

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

JC979 U.S. PTO
10/087573
02/28/02

In re the application of:

SCHETTERS Theodorus P.M.

CARCY Bernard P.D.

DRACULOVSKI Pascal E.

GOEENFLOT Andre F.

Serial Number: To be assigned Group Art Unit: To be assigned
Filed: concurrently herewith Examiner: To be assigned
For: BABESIA CANIS VACCINE

Corresponding to: EP01200316.5, Filed 06 March, 2001

CLAIM TO PRIORITY UNDER 35 USC 119

Assistant Commissioner of Patents
Washington, D.C. 20231

February 28, 2002

Sir:

The benefit of the filing date of the following prior foreign application is hereby requested for the above-identified application, and the priority provided in 35 USC 119 is hereby claimed:

European Patent Application No. 01200816.5, filed March 6, 2001

In support of this claim, the requisite certified copy of said original foreign application is filed herewith along with a verified English translation thereof.

It is requested that the file of this application be marked to indicate that the Applicant has complied with the requirements of 35 USC 119 and that the Patent and Trademark Office kindly

100

101

102

103

104

105

106

107

108

109

110

111

112

113

114

115

116

117

118

119

120

121

122

123

124

125

126

127

128

129

130

131

132

133

134

135

136

137

138

139

140

141

142

143

144

145

146

147

148

149

150

151

152

153

154

155

156

157

158

159

160

161

162

163

164

165

166

167

168

169

170

171

172

173

174

175

176

177

178

179

180

181

182

183

184

185

186

187

188

189

190

191

192

193

194

195

196

197


198

199

200

In the event any fees are required with this paper, please
charge our Deposit Account No. 02-2334.

Respectfully submitted,

A handwritten signature in black ink, appearing to read "William P. Ramey III", written over a horizontal line.

William P Ramey III
Patent Counsel
Registration Number 44,295

Attorney Docket No. I-2001.004 US

Intervet Inc Patent Department
Akzo Nobel
405 State Street
P.O. Box 318
Millsboro, DE 19966
Tel: (302) 933-4034
Fax: (302) 933-4013

WPR:kmm



Europäisches
Patentamt

European
Patent Office

Office européen
des brevets

JC979 U.S. PTO

10/087573



BEST AVAILABLE COPY

Bescheinigung

Certificate

Attestation

Die angehefteten Unterla-
gen stimmen mit der
ursprünglich eingereichten
Fassung der auf dem näch-
sten Blatt bezeichneten
europäischen Patentanmel-
dung überein.

The attached documents
are exact copies of the
European patent application
described on the following
page, as originally filed.

Les documents fixés à
cette attestation sont
conformes à la version
initialement déposée de
la demande de brevet
européen spécifiée à la
page suivante.

Patentanmeldung Nr. Patent application No. Demande de brevet n°

01200816.5

Der Präsident des Europäischen Patentamts;
Im Auftrag

For the President of the European Patent Office

Le Président de l'Office européen des brevets
p.o.

DEN HAAG, DEN
THE HAGUE,
LA HAYE, LE

11/01/02





Europäisches
Patentamt

European
Patent Office

Office européen
des brevets

Blatt 2 der Bescheinigung
Sheet 2 of the certificate
Page 2 de l'attestation

Anmeldung Nr.:
Application no.:
Demande n°: 01200816.5

Anmeldetag:
Date of filing:
Date de dépôt: 06/03/01

Anmelder:
Applicant(s):
Demandeur(s):
Akzo Nobel N.V.
6824 BM Arnhem
NETHERLANDS

Bezeichnung der Erfindung
Title of the invention:
Titre de l'invention:
Babesia canis vaccine

In Anspruch genommene Priorität(en) / Priority(ies) claimed / Priorité(s) revendiquée(s)

Staat
State:
Pays

Tag
Date:
Date

Aktenzeichen:
File no.
Numéro de dépôt

Internationale Patentklassifikation
International Patent classification
Classification internationale des brevets

/

Am Anmeldetag benannte Vertragsstaaten
Contracting states designated at date of filing: AT/BE/CH/CY/DE/DK/ES/FI/FR/GB/GR/IE/IT/LI/LU/MC/NL/PT/SE/TR
Etats contractants désignés lors du dépôt

Bemerkungen
Remarks

100

101

102

103

104

105

106

107

108

109

110

111

112

113

114

115

116

117

118

119

120

121

122

123

124

125

126

127

128

129

130

131

132

133

134

135

136

137

138

139

140

141

142

143

144

145

146

147

148

149

150

151

152

153

154

155

156

157

158

159

160

161

162

163

164

165

166

167

168

169

170

171

172

173

174

175

176

177

178

179

180

181

182

183

184

185

186

187

188

189

190

191

192

193

194

195

196

197

198

199

200

201

202

203

204

205

206

207

208

209

210

211

212

213

214

215

216

217

218

219

220

221

222

223

224

225

226

227

228

229

230

231

232

233

234

235

236

237

238

239

240

241

242

243

244

245

246

247

248

249

250

251

252

253

254

255

256

257

258

259

260

261

262

263

264

265

266

267

268

269

270

271

272

273

274

275

276

277

278

279

280

281

282

283

284

285

286

287

288

289

290

291

292

293

294

295

296

297

298

299

300

301

302

303

304

305

306

307

308

309

310

311

312

313

314

315

316

317

318

319

320

321

322

323

324

325

326

327

328

329

330

331

332

333

334

335

336

337

338

339

340

341

342

343

344

345

346

347

348

349

350

351

352

353

354

355

356

357

358

359

360

361

362

363

364

365

366

367

368

369

370

371

372

373

374

375

376

377

378

379

380

381

382

383

384

385

386

387

388

389

390

391

392

393

394

395

396

397

398

399

400

401

402

403

404

405

406

407

408

409

410

411

412

413

414

415

416

417

418

419

420

421

422

423

424

425

426

427

428

429

430

431

432

433

434

435

436

437

438

439

440

441

442

443

444

445

446

447

448

449

450

451

452

453

454

455

456

457

458

459

460

461

462

463

464

465

466

467

468

469

470

471

472

473

474

475

476

477

478

479

480

481

482

483

484

485

486

487

488

489

490

491

492

493

494

495

496

497

498

499

500

501

502

503

504

505

506

507

508

509

510

511

512

513

514

515

516

517

518

519

520

521

522

523

524

525

526

527

528

529

530

531

532

533

534

535

536

537

538

539

540

541

542

543

544

545

546

547

548

549

550

551

552

553

554

555

556

557

558

559

560

561

562

563

564

565

566

567

568

569

570

571

572

573

574

575

576

577

578

579

580

581

582

583

584

585

586

587

588

589

590

591

592

593

594

595

596

597

598

599

600

601

602

603

604

605

606

607

608

609

610

611

612

613

614

615

616

617

618

619

620

621

622

623

624

625

626

627

628

629

630

631

632

633

634

635

636

637

638

639

640

641

642

643

644

645

646

647

648

649

650

651

652

653

654

655

656

657

658

659

660

661

662

663

664

665

666

667

668

669

670

671

672

673

674

675

676

677

678

679

680

681

682

683

684

685

686

687

688

689

690

691

692

693

694

695

696

697

698

699

700

701

702

703

704

705

706

707

708

709

710

711

712

713

714

715

716

717

718

719

720

721

722

723

724

725

726

727

728

729

730

731

732

733

734

735

736

737

738

739

740

741

742

743

744

745

746

747

748

749

750

751

752

753

754

755

756

757

758

759

760

761

762

763

764

765

766

767

768

769

770

771

772

773

774

775

776

777

778

779

780

781

782

783

784

785

786

787

788

789

790

791

792

793

794

795

796

797

798

799

800

801

802

803

804

805

806

807

808

809

810

811

812

813

814

815

816

817

818

819

820

821

822

823

824

825

826

827

828

829

830

831

832

833

834

835

836

837

838

839

840

841

842

843

844

845

846

847

848

849

850

851

852

853

854

855

856

857

858

859

860

861

862

863

864

865

866

867

868

869

870

871

872

873

874

875

876

877

878

879

880

881

882

883

884

885

886

887

888

889

890

891

892

893

894

895

896

897

898

899

900

901

902

903

904

905

906

907

908

909

910

911

912

913

914

915

916

917

918

919

920

921

922

923

924

925

926

927

928

929

930

931

932

933

934

935

936

937

938

939

940

941

942

943

944

945

946

947

948

949

950

951

952

953

954

955

956

957

958

959

960

961

962

963

964

965

966

967

968

969

970

971

972

973

974

975

976

977

978

979

980

981

982

983

984

985

986

987

988

989

990

991

992

993

994

995

996

997

998

999

1000

Babesia canis vaccine.

06. 03. 2001

The present invention relates to nucleic acid sequences encoding ⁽³⁸⁾ novel *Babesia canis* associated proteins, to cDNA fragments, recombinant DNA molecules and live
 5 recombinant carriers comprising these sequences and to host cells comprising such nucleic acid sequences, cDNA fragments, recombinant DNA molecules and live recombinant carriers. Furthermore, the invention relates to proteins encoded by these nucleotide sequences, to vaccines for combating *Babesia canis* infections and methods for the preparation thereof, to *Babesia canis* associated antigenic material for
 10 use in vaccines and to the use of *Babesia canis* associated antigenic material in the manufacture of vaccines. Also, the invention relates to diagnostic tools for the detection of *Babesia canis* associated nucleic acid sequences, for the detection of *Babesia canis* associated antigens and for the detection of antibodies against *Babesia canis* associated antigenic material.

15 Babesiosis, like malaria, is a disease which has a focal character. The reason for this is that the pathogen is transmitted by ticks that feed on a certain reservoir of parasites present in the vertebrate population. Only where ticks are present, babesiosis can occur. On balance, particularly in indigenous animals, the parasite coexists with the
 20 host without causing significant disease. In many cases babesiosis is a problem because of man's activities through inbreeding of genetic traits and/or transporting animals to unfamiliar environments where babesiosis is endemic (Callow, L.L and Dalglish, R.J., 1982).

In dogs the disease is caused by *B. canis* (transmitted by Dermacentor ticks), *B. rossi*
 25 (transmitted by Haemaphysalis ticks), *B. vogeli* (transmitted by Rhipicephalus ticks) and *Babesia gibsoni* (transmitted by Haemaphysalis and Rhipicephalus ticks). Signs of disease in naturally acquired babesiosis usually begin 7-21 days after infection. These symptoms include: fever, anorexia, depression, anaemia, haemoglobinuria and rapidly developing weakness. Increased lacrimation, salivation
 30 and muscle tremor commonly occur. Nervous signs may develop in terminal

clinical cases resembling disseminated intravascular coagulation (DIC) have been reported in acute *B. canis* infections. Thrombosis is not common, but small hyaline thrombi, connected with megakaryocytes have been described. It appears that these

coagulation disturbances lead to increased erythrocyte-stickiness. As a result the blood passage through the microvasculature is hampered, resulting in congestion of internal organs and decreased packed cell volumes (PCV). This might impair the oxygen supply to certain tissues and subsequently lead to tissue damage as a result of anoxia. Evidence from congestion in *B. canis* infections comes from experiments in which dogs were chemotherapeutically treated. Some of these animals restore the packed cell volume in two days from 25-35%, which is associated with shrinkage of the spleen to normal measures.

The four species of *Babesia* differ with respect to their pathogenicity. The

North-African *B. vogeli* strains provoke only mild disease, which usually does not require treatment (like the Australian *B. canis* strains). European *B. canis* strains are more pathogenic than the North-African parasite. In our experiments with the *B. canis* A strain, approximately half of the animals required chemotherapeutic treatment after infection. The South-African *B. rossi* strains are most pathogenic, a feature observed already very early. Using the South-African *B. rossi* strain naive dogs developed progressing disease characterised by exponential parasite growth. In contrast, parasitaemia in dogs infected with European *B. canis* usually is limited. In the latter, congestion appears to be the main pathological feature.

At the moment a vaccine for *Babesia* is prepared from the supernatant of a culture of a strain of *Babesia canis*, as is described in US Patent 4,777,036. However, such a vaccine contains only antigens of the strain of this particular species of *B. canis*. It appears that such a vaccine gives in general little protection against infections with (wild type) *B. canis* (Lepetit, C., Piroplasmose canine et vaccination Pirodog, Doctoral Thesis, Univ. of Nantes, 1988). The author of the thesis gives three possible causes of this failure of protective immunisation:

1. the sensitivity of the dogs with respect to vaccination;
2. the type of immune effector mechanism induced by vaccination;
3. the antigenic diversity of *Babesia*.

Vaccines containing whole, attenuated *Babesia* parasites always harbour the danger that the parasites again become virulent and thus spread the disease instead of curing it. Therefore, the use of subunit vaccines is preferred. Thus there remains the desire for a vaccine which is not infective, which can be easily produced and which yet can

give protection against *Babesia canis* infection, preferably against all *Babesia canis* strains.

5 It is i.a. an objective of the present invention to provide *Babesia canis* related proteins which are able, each separately, or when combined, to induce protection against *Babesia canis* infection in dogs. It was surprisingly found now, that two such novel proteins exist that are specifically associated with *Babesia canis*: they are not found in the *Babesia* species *B. rossi*.

10 By referring to the novel proteins as associated with *Babesia canis* it is meant that these proteins are found to be present in the *Babesia canis* parasite. It does not mean that the genetic information is found on a chromosome of the parasite. The genetic information for the novel proteins is found in the parasite, but it can be present as extra-chromosomal information.

15 The novel proteins will be referred to as the 15 kD protein (also BcVir15) and the 32 kD protein (also BcVir32). The amino acid sequence of the 15 kD-protein is presented in sequence identifier SEQ ID NO: 2. The cDNA encoding this protein has been sequenced and its nucleic acid sequence is shown in sequence identifier SEQ ID NO:
20 1.

The amino acid sequence of the 32 kD protein is presented in sequence identifier SEQ ID NO: 4. The cDNA encoding this protein has been sequenced and its nucleic acid sequence is shown in sequence identifier SEQ ID NO: 3.

25 It is well-known in the art, that many different nucleic acid sequences can encode one and the same protein. This phenomenon is commonly known as wobble in the second and especially the third base of each triplet encoding an amino acid. This phenomenon can result in a heterology of about 30% for two different nucleic acid sequences still
30 encoding the same protein. Therefore, two nucleic acid sequences having a sequence

One embodiment of the invention relates to a nucleic acid sequence encoding a 15 kD *Babesia canis* associated protein or an immunogenic fragment of that protein, said

protein or immunogenic fragment thereof having at least 80 %, preferably 90 %, more preferably 95 % homology with the amino acid sequence as depicted in SEQ ID NO:

2. Even more preferred is a homology level of 98% or even 100%.

- 5 Another embodiment of the invention relates to a nucleic acid sequence encoding a 32 kD *Babesia canis* associated protein or an immunogenic fragment of that protein, said protein or immunogenic fragment thereof having at least 80 %, preferably 90 %, more preferably 95 % homology with the amino acid sequence as depicted in SEQ ID NO:
4. Even more preferred is a homology level of 98% or even 100%.

10

Needless to say that a nucleic acid sequence encoding a protein according to the invention is understood to apply equally to the RNA encoding the protein and to cDNA and DNA carrying the genetic code of this RNA. As a consequence, where a certain nucleic acid sequence or part thereof is said to have a certain level of

- 15 homology with e.g. the nucleic acid sequence of SEQ ID NO: 1, this level of homology equally goes for cDNA and the RNA used as a template for making the cDNA, as well as vice versa to (c)DNA and the RNA transcribed from that DNA.

- Parts of the said nucleic acid sequences that encode an immunogenic fragment of the
- 20 proteins to which the invention relates can be very small. Immunogenic fragments of e.g. 4 amino acid length are known. Such immunogenic fragments are encoded by a nucleic acid sequence of 12 nucleotides. Such very short amino acid sequences are however not immunogenic as such: they need to be coupled to a carrier, such as KHL, as known in the art. Longer fragments can be immunogenic as such. Therefore, the
- 25 parts of the said nucleic acid sequences that encode an immunogenic fragment comprise at least 12 nucleotides. Parts comprising more than 15, more preferably more than 18, 20 25 or even 30 nucleotides are even more preferred in that order of preference.

- 30 Since the present invention discloses nucleic acid sequences encoding novel *Babesia canis* associated 15 kD and 32 kD proteins, it is now for the first time possible to obtain these proteins in sufficient quantities. This can e.g. be done by using expression systems to express the genes encoding the proteins.

Therefore, in another embodiment, the invention relates to a cDNA fragment comprising a nucleic acid sequence according to the invention. A cDNA is a copy DNA that has been made, e.g. by reverse transcriptase, of an RNA encoding a protein according to the invention. Such cDNA fragments in their shortest form are a DNA
 5 copy of a part of a nucleotide sequence according to the invention. They can however also comprise flanking nucleic acid sequences not naturally present upstream and downstream of the coding sequence.

An essential requirement for the expression of cDNA is an adequate promoter
 10 functionally linked to the cDNA, so that the cDNA is under the control of the promoter. It is obvious to those skilled in the art that the choice of a promoter extends to any eukaryotic, prokaryotic or viral promoter capable of directing gene transcription in cells used as host cells for protein expression.

Therefore, another embodiment of the invention relates to a recombinant DNA
 15 molecule comprising a cDNA fragment or a nucleic acid sequence according to the invention that is placed under the control of a functionally linked promoter. This can be obtained by means of e.g. standard molecular biology techniques.

(Maniatis/Sambrook (Sambrook, J. Molecular cloning: a laboratory manual, 1989. ISBN 0-87969-309-6).

20 Functionally linked promoters are promoters that are capable of controlling the transcription of the nucleic acid sequences to which they are linked.

Such a promoter can be a *Babesia* promoter, provided that that promoter is functional in the cell used for expression. It can also be a heterologous promoter. When the host cells are bacteria, useful expression control sequences which may be used include the
 25 Trp promoter and operator (Goeddel, et al., Nucl. Acids Res., 8, 4057, 1980); the lac promoter and operator (Chang, et al., Nature, 275, 615, 1978); the outer membrane protein promoter (Nakamura, K. and Inouge, M., EMBO J., 1, 771-775, 1982); the bacteriophage lambda promoters and operators (Remaut, E. et al., Nucl. Acids Res., 11, 4677-4688, 1983); the α -amylase (*B. subtilis*) promoter and operator, termination
 30 sequences and other expression enhancement and control sequences compatible with

When the host cell is yeast, useful expression control sequences include, for

example, mating factor. For insect cells the polyhedrin or p10 promoters of baculoviruses can

be used (Smith, G.E. et al., Mol. Cell. Biol. 3, 2156-65, 1983). When the host cell is of mammalian origin illustrative useful expression control sequences include the SV-40 promoter (Berman, P.W. et al., Science, 222, 524-527, 1983) or the metallothionein promoter (Brinster, R.L., Nature, 296, 39-42, 1982) or a heat shock promoter (Voellmy et al., Proc. Natl. Acad. Sci. USA, 82, 4949-53, 1985).

Bacterial, yeast, fungal, insect and mammalian cell expression systems are very frequently used systems. Such systems are well-known in the art and generally available, e.g. commercially through Clontech Laboratories, Inc. 4030 Fabian Way, Palo Alto, California 94303-4607, USA. Next to these expression systems, parasite-based expression systems are very attractive expression systems. Such systems are e.g. described in the French Patent Application with Publication number 2 714 074, and in US NTIS Publication No US 08/043109 (Hoffman, S. and Rogers, W.: Public. Date 1 December 1993).

Another embodiment of the invention relates to a Live Recombinant Carrier (LRC) comprising a cDNA fragment according to the invention or a recombinant DNA molecule according to the invention. Such carriers are e.g. bacteria, parasites and viruses. These LRCs are micro-organisms or viruses in which additional genetic information, in this case a nucleic acid sequence encoding the 15 kD or 32 kD protein or an immunogenic fragment thereof according to the invention has been cloned. Animals infected with such LRCs will produce an immunogenic response not only against the immunogens of the carrier, but also against the immunogenic parts of the protein(s) for which the genetic code is additionally cloned into the LRC, e.g. the 15 kD or 32 kD cDNA.

As an example of bacterial LRCs, attenuated *Salmonella* strains known in the art can attractively be used.

Live recombinant carrier parasites have i.a. been described by Vermeulen, A. N. (Int. Journ. Parasitol. 28: 1121-1130 (1998)).

Also, LRC viruses may be used as a way of transporting the nucleic acid sequence into a target cell. Live recombinant carrier viruses are also called vector viruses. Viruses often used as vectors are Vaccinia viruses (Panicali et al; Proc. Natl. Acad. Sci. USA, 79: 4927 (1982), Herpesviruses (E.P.A. 0473210A2), and Retroviruses (Valerio, D. et al; in Baum, S.J., Dicke, K.A., Lotzova, E. and Pluznik, D.H. (Eds.),

The technique of *in vivo* homologous recombination, well-known in the art, can be used to introduce a recombinant nucleic acid sequence into the genome of a bacterium, parasite or virus of choice, capable of inducing expression of the inserted nucleic acid sequence according to the invention in the host animal.

Finally another embodiment of the invention relates to a host cell comprising a nucleic acid sequence according to the invention, a cDNA fragment or a recombinant DNA molecule according to the invention or a live recombinant carrier according to the invention. A host cell may be a cell of bacterial origin, e.g. *Escherichia coli*, *Bacillus subtilis* and *Lactobacillus* species, in combination with bacteria-based plasmids as pBR322, or bacterial expression vectors as pGEX, or with bacteriophages. The host cell may also be of eukaryotic origin, e.g. yeast-cells in combination with yeast-specific vector molecules, or higher eukaryotic cells like insect cells (Luckow et al; Bio-technology 6: 47-55 (1988)) in combination with vectors or recombinant baculoviruses, plant cells in combination with e.g. Ti-plasmid based vectors or plant viral vectors (Barton, K.A. et al; Cell 32: 1033 (1983)), mammalian cells like Hela cells, Chinese Hamster Ovary cells (CHO) or Crandell Feline Kidney-cells, also with appropriate vectors or recombinant viruses.

Another embodiment of the invention relates to the novel *Babesia canis* associated proteins: the 15 kD and 32 kD protein and to immunogenic fragments thereof, according to the invention.

The concept of immunogenic fragments will be defined below.

It will be understood that, for the particular proteins embraced herein, natural variations can exist between the proteins associated with individual *Babesia canis* strains. These variations may be demonstrated by (an) amino acid difference(s) in the overall sequence or by deletions, substitutions, insertions, inversions or additions of

Other biological and immunological activities, have been described, e.g., in a summary article in "The Proteins" Academic Press New York (1979). Amino acid replacements between related amino acids or replacements which have occurred frequently in

evolution are, inter alia, Ser/Ala, Ser/Gly, Asp/Gly, Asp/Asn, Ile/Val (see Dayhof, M.D., Atlas of protein sequence and structure, Nat. Biomed. Res. Found., Washington D.C., 1978, vol. 5, suppl. 3). Other amino acid substitutions include Asp/Glu, Thr/Ser, Ala/Gly, Ala/Thr, Ser/Asn, Ala/Val, Thr/Phe, Ala/Pro, Lys/Arg, Leu/Ile, Leu/Val and
 5 Ala/Glu. Based on this information, Lipman and Pearson developed a method for rapid and sensitive protein comparison (Science, 227, 1435-1441, 1985) and determining the functional similarity between homologous proteins. Such amino acid substitutions of the exemplary embodiments of this invention, as well as variations having deletions and/or insertions are within the scope of the invention as long as the
 10 resulting proteins retain immune reactivity.

This explains why *Babesia canis* associated proteins according to the invention, when isolated from different field isolates, may have homology levels of about 80%, while still representing the same protein with the same immunological characteristics, i.e. the capability of inducing an immunological response against a *Babesia canis*
 15 infection.

Those variations in the amino acid sequence of a certain protein according to the invention that still provide a protein capable of inducing an immune response against infection with *Babesia canis* or at least against the clinical manifestations of the infection are considered as "not essentially influencing the immunogenicity".
 20

Therefore, one form of this embodiment relates i.a. to a *Babesia canis* associated protein that has a molecular weight of 15 kD and comprises an amino acid sequence that is at least 80 % homologous to the amino acid sequence as depicted in SEQ ID NO: 2 and to an immunogenic fragment of said protein.
 25

In a preferred form, the embodiment relates to such a *Babesia canis* associated protein that has a sequence homology of at least 85 %, preferably 90 %, more preferably 95 % homology to the amino acid sequence as depicted in SEQ ID NO: 2 and to an immunogenic fragment of such a protein.

30 Even more preferred is a homology level of 98% or even 100%.

Another embodiment relates i.a. to a *Babesia canis* associated protein that has a molecular weight of 32 kD and comprises an amino acid sequence that is at least 80 % homologous to the amino acid sequence as depicted in SEQ ID NO: 4 and to an

immunogenic fragment of said protein.

A preferred form of this embodiment relates to such a *Babesia canis* associated protein that has a sequence homology of at least 85 %, preferably 90 %, more preferably 95 % homology to the amino acid sequence as depicted in SEQ ID NO: 4 and to an immunogenic fragment of such a protein.
Even more preferred is a homology level of 98% or even 100%.

The level of protein homology can be determined with the computer program "BLAST 2 SEQUENCES" by selecting sub-program: "BLASTP", that can be found at www.ncbi.nlm.nih.gov/blast/bl2seq/bl2.html.

A reference for this program is Tatjana A. Tatusova, Thomas L. Madden FEMS Microbiol. Letters 174: 247-250 (1999). Matrix used: "blosum62". Parameters used are the default parameters: Open gap: 11. Extension gap: 1. Gap x_dropoff: 50.

When a protein is used for e.g. vaccination purposes or for raising antibodies, it is however not necessary to use the whole protein. It is also possible to use a fragment of that protein that is capable, as such or coupled to a carrier such as e.g. KLH, of inducing an immune response against that protein, a so-called immunogenic fragment.
An "immunogenic fragment" is understood to be a fragment of the full-length protein that still has retained its capability to induce an immune response in the host, i.e. comprises a B- or T-cell epitope. At this moment, a variety of techniques is available to easily identify cDNA fragments encoding antigenic fragments (determinants). The method described by Geysen et al (Patent Application WO 84/03564, Patent Application WO 86/06487, US Patent NR. 4,833,092, Proc. Natl Acad. Sci. 81: 3998-4002 (1984), J. Imm. Meth. 102, 259-274 (1987)), the so-called PEPSCAN method is an easy to perform, quick and well-established method for the detection of epitopes, the immunologically important regions of the protein. The method is used world-wide and as such well-known to man skilled in the art. This (empirical) method is especially suitable for the detection of B-cell epitopes. Also, given the sequence of the

fragments as the immunogenic protein, fragments capable of inducing an immune response and/or structural agreement with epitopes that are now known. The determination of these regions is based on a combination of the hydrophilicity criteria according to

Hopp and Woods (Proc. Natl. Acad. Sci. 78: 38248-3828 (1981)), and the secondary structure aspects according to Chou and Fasman (Advances in Enzymology 47: 45-148 (1987) and US Patent 4,554,101). T-cell epitopes can likewise be predicted from the sequence by computer with the aid of Berzofsky's amphiphilicity criterion (Science 235, 1059-1062 (1987) and US Patent application NTIS US 07/005,885). A condensed overview is found in: Shan Lu on common principles: Tibtech 9: 238-242 (1991), Good et al on Malaria epitopes; Science 235: 1059-1062 (1987), Lu for a review; Vaccine 10: 3-7 (1992), Berzowsky for HIV-epitopes; The FASEB Journal 5:2412-2418 (1991).

10

Thus, another embodiment of the invention relates to a vaccine for combating *Babesia canis* infection, that comprises a protein or an immunogenic fragment thereof according to the invention as described above together with a pharmaceutically acceptable carrier.

15

One way of making a vaccine according to the invention comprises growing the *Babesia canis* parasite, and purification of the 15 kD and 32 kD proteins. This is however a very time-consuming way of making the vaccine. It would thus be less practical to make such vaccines on an industrial scale.

20

It is one of the merits of the present invention that it provides for the first time the nucleic acid sequences encoding the 15 kD and 32 kD proteins. It is therefore much more convenient to use the expression products of the cDNA encoding these proteins or immunogenic fragments thereof according to the invention in vaccines.

25

Vaccines based upon the expression products of these cDNAs can easily be made by admixing one or both proteins according to the invention or immunogenic fragments thereof according to the invention with a pharmaceutically acceptable carrier as described below. It is also possible to directly admix a host cell in which one or both proteins according to the invention or immunogenic fragments thereof according to the invention have been expressed and a pharmaceutically acceptable carrier.

30

Alternatively, a vaccine according to the invention can comprise live recombinant carriers as described above, capable of expressing the proteins according to the

invention or immunogenic fragments thereof according to the invention. Such vaccines, e.g. based upon a bacterial (e.g. *Salmonella*) carrier, a parasitic or a viral carrier have the advantage over subunit vaccines that they better mimic the natural way of infection of *Babesia canis*. Moreover, their self-propagation is an advantage since only low amounts of the recombinant carrier are necessary for immunisation. Thus, still another embodiment relates to a vaccine for combating *Babesia canis* infection, that comprises a live recombinant carrier according to the invention and a pharmaceutically acceptable carrier.

10 All vaccines described above contribute to active vaccination, i.e. the host's immune system is triggered by one or more proteins according to the invention or immunogenic fragments thereof, to make antibodies against these proteins or fragments or to expand T-cell populations with receptors that can be specifically triggered by epitopes associated with these proteins.

15 Alternatively, such antibodies can be raised in e.g. rabbits or can be obtained from antibody-producing cell lines as described below. Such antibodies can then be administered to the host animal. This method of vaccination, passive vaccination, is the vaccination of choice when an animal is already infected, and there is no time to allow the natural immune response to be triggered. It is also the preferred method for
20 vaccinating immune-compromised animals. Administered antibodies against *Babesia canis* can in these cases bind directly to the proteins according to the invention and to parasites and cells exposing these proteins. This has the advantage that it decreases or stops *Babesia canis* growth.

Therefore, one other form of this embodiment of the invention relates to a vaccine for
25 combating *Babesia canis* infection comprising antibodies against a *Babesia canis* associated protein according to the invention or an immunogenic fragment of those proteins, and a pharmaceutically acceptable carrier.

Another embodiment relates to a method for the preparation of a vaccine according to
30 the invention that comprises the admixing of antibodies according to the invention

Vaccines can also be based upon host cells as described above, that comprise the nucleic acid sequences, cDNA fragments, recombinant DNA molecules or live recombinant carriers according to the invention.

Therefore, another embodiment of the present invention relates to a vaccine for
 5 combating *Babesia canis* infection, that comprises a host cell according to the invention and a pharmaceutically acceptable carrier.

An alternative and efficient way of vaccination is direct vaccination with DNA encoding the relevant antigen. Direct vaccination with DNA encoding proteins has
 10 been successful for many different proteins (As reviewed in e.g. Donnelly et al., The Immunologist 2: 20-26 (1993)).

This way of vaccination is also attractive for the vaccination of dogs against *Babesia canis* infection.

Therefore, still other forms of this embodiment of the invention relate to a vaccine
 15 comprising a nucleic acid sequence according to the invention and a pharmaceutically acceptable carrier, and to a vaccine comprising a cDNA fragment according to the invention and a pharmaceutically acceptable carrier.

Still another form of this embodiment relates to a vaccine comprising a recombinant DNA molecule according to the invention and a pharmaceutically acceptable carrier.
 20 DNA vaccines can easily be administered through intradermal application e.g. using a needle-less injector. This way of administration delivers the DNA directly into the cells of the animal to be vaccinated. Amounts of DNA in the microgram range between 1 and 100 µg provide very good results.

25 In a preferred embodiment, the vaccine according to the present invention comprises an additional antigen derived from a micro-organism or virus pathogenic to dogs, or genetic information encoding such an antigen.

Such dog pathogenic organisms and viruses are preferably selected from the group of *Ehrlichia canis*, *Babesia gibsoni*, *vogeli*, *rossi*, *Leishmania donovani*-complex,
 30 Canine parvovirus, Canine distempervirus, *Leptospira interrogans serovar canicola*, *icterohaemorrhagiae*, *pomona*, *grippotyphosa*, *bratislava*, Canine hepatitisvirus, Canine parainfluenzavirus, rabies virus, *Hepatozoon canis* and *Borrelia burgdorferi*.

All vaccines according to the present invention comprise a pharmaceutically acceptable carrier. A pharmaceutically acceptable carrier can be e.g. sterile water or a sterile physiological salt solution. In a more complex form the carrier can e.g. be a buffer.

5

Methods for the preparation of a vaccine comprise the admixing of a nucleic acid sequence, a cDNA fragment, a recombinant DNA molecule or a live recombinant carrier according to the invention, a host cell according to the invention, or a protein according to the invention or an immunogenic fragment thereof and a pharmaceutically acceptable carrier.

10

Thus, another embodiment of the present invention relates to a method for the preparation of a vaccine according to the invention, that comprises the admixing of a nucleic acid sequence according to the invention, a cDNA fragment the invention, a recombinant DNA molecule according to the invention, a live recombinant carrier according to the invention or a host cell according to the invention or a protein according to the invention and a pharmaceutically acceptable carrier.

15

Vaccines according to the present invention may in a preferred presentation also contain an adjuvant. Adjuvants in general comprise substances that boost the immune response of the host in a non-specific manner. A number of different adjuvants are known in the art. Examples of adjuvants are Freund's Complete and Incomplete adjuvant, vitamin E, non-ionic block polymers, muramyldipeptides, saponins such as Quill A(R), mineral oil e.g. Bayol(R) or Markol(R), vegetable oil, and Carbopol(R) (a homopolymer), or Diluvac(R) Forte.

20

The vaccine may also comprise a so-called "vehicle". A vehicle is a compound to which the protein adheres, without being covalently bound to it. Often used vehicle compounds are e.g. aluminium hydroxide, -phosphate or -oxide, silica, Kaolin, and Bentonite.

25

A special form of such a vehicle, in which the antigen is partially embedded in the vehicle, is the so called ISCOM (EP 109 942, EP 180 564, EP 242 380)

30

with emulsifiers, e.g. Span or Tween

Often, the vaccine is mixed with stabilisers, e.g. to protect degradation-prone proteins from being degraded, to enhance the shelf-life of the vaccine, or to improve freeze-drying efficiency. Useful stabilisers are i.a. SPGA (Bovarnik et al; J. Bacteriology 59: 509 (1950)), carbohydrates e.g. sorbitol, mannitol, trehalose, starch, sucrose, dextran or glucose, proteins such as albumin or casein or degradation products thereof, and buffers, such as alkali metal phosphates.

In addition, the vaccine may be suspended in a physiologically acceptable diluent. It goes without saying, that other ways of adjuvating, adding vehicle compounds or diluents, emulsifying or stabilising a protein are also embodied in the present invention.

Thus, a preferred form of this embodiment relates to a vaccine for combating *Babesia canis* infections according to the invention that comprises an adjuvant.

Vaccines according to the invention can very suitably be administered in amounts ranging between 1 and 100 micrograms, although smaller doses can in principle be used. A dose exceeding 100 micrograms will, although immunologically very suitable, be less attractive for commercial reasons.

Vaccines based upon live attenuated recombinant carriers, such as the LRC-viruses, parasites and bacteria described above can be administered in much lower doses, because they multiply themselves during the infection. Therefore, suitable amounts would range between 10^3 and 10^9 CFU/PFU for respectively bacteria, parasites and viruses.

Many ways of administration can be applied. Systemic application is a suitable way of administration, e.g. by intramuscular application of the vaccine. If this route is followed, standard procedures known in the art for systemic application are well-suited. Intradermal, sub-cutaneous and intracutaneous ways of administration are also attractive.

Still another embodiment of the present invention relates to a *Babesia canis* associated protein according to the invention for use in a vaccine.

Still another embodiment of the present invention relates to the use of a *Babesia canis* associated protein according to the invention for the manufacturing of a vaccine for combating *Babesia canis* infections.

- 5 From a point of view of protection against disease, a quick and correct diagnosis of *Babesia canis* infection is important.

Therefore it is another objective of this invention to provide diagnostic tools suitable for the detection of *Babesia canis* infection.

- 10 A diagnostic test for the detection of *Babesia canis* is e.g. based upon the reaction of RNA isolated from the animal to be tested, with specific probes or it is e.g. a RT-PCR test based upon the coding sequences for the 15 kD or the 32 kD proteins or based upon nucleic acid sequences that are complementary to those coding sequences. If nucleic acid molecules specific for the *Babesia canis* associated proteins according to
- 15 the invention are present in the animal, these will e.g. specifically bind to specific PCR-primers and will subsequently become amplified in RT-PCR-reaction. The PCR-reaction product can then easily be detected in DNA gel electrophoresis. RT-PCR reactions are well-known in the art (see reference below). The nucleic acid molecules can most easily be isolated from (the parasites present in) erythrocytes of the animal
- 20 to be tested. Standard PCR-textbooks give methods for determining the length of the primers for selective PCR-reactions with nucleic acid molecules specific for the *Babesia canis* associated proteins according to the invention. Primers with a nucleotide sequence of at least 12 nucleotides are frequently used, but primers of more than 15, more preferably 18 nucleotides are somewhat more selective.
- 25 Especially primers with a length of at least 20, preferably at least 30 nucleotides are very generally applicable. PCR-techniques are extensively described in (Dieffenbach & Drexler; PCR primers, a laboratory manual. ISBN 0-87969-447-5 (1995)). Nucleic acid molecules encoding a *Babesia canis* associated protein or parts of those nucleic acid molecules having a length of at least 12, preferably 15, more preferably
- 30 18, even more preferably 20, 22, 25, 30, 35 or 40 nucleotides in that order of

nucleic acid sequence that is complementary to nucleic acid sequence as depicted in SEQ ID NO: 1 or 3 are therefore also part of the invention. Such nucleic acid

molecules can e.g. be used as primers in (RT-)PCR-reactions in order to enhance the amount of nucleic acid that encodes the proteins according to the invention. This allows the quick amplification of specific nucleotide sequences for use as a diagnostic tool for e.g. the detection of *Babesia* in tissue as indicated above.

5

The level of nucleotide homology can be determined with the computer program "BLAST 2 SEQUENCES" by selecting sub-program: "BLASTN" that can be found at www.ncbi.nlm.nih.gov/blast/bl2seq/bl2.html.

A reference for this program is Tatiana A. Tatusova, Thomas L. Madden FEMS Microbiol. Letters 174: 247-250 (1999). Parameters used are the default parameters: Reward for a match: +1. Penalty for a mismatch: -2. Open gap: 5. Extension gap: 2. Gap x_dropoff: 50.

15

Another nuclei acid-based test is based upon growth of parasite material obtained from erythrocytes, followed by classical RNA purification followed by cDNA synthesis and/or classical hybridisation with radioactively or colour labelled 15 kD or 32 kD protein-specific cDNA-fragments. Both PCR-reactions and hybridisation reactions are well-known in the art and are i.a. described in Maniatis/Sambrook (Sambrook, J. et al. Molecular cloning: a laboratory manual. ISBN 0-87969-309-6).

20

Thus, another embodiment of the invention relates to a diagnostic test for the detection of *Babesia canis* associated RNA characterised in that the test comprises a nucleic acid sequence that is at least 70 % homologous to the nucleic acid sequence as depicted in SEQ ID NO: 1 or 3 or a nucleotide sequence that is complementary to that nucleic acid sequence, or a fragment thereof having a length of at least 12, preferably 15, more preferably 18 nucleotides.

25

A diagnostic test for the detection of antibodies against the *Babesia canis* associated proteins in e.g. sera, tissue or body fluids can be e.g. a simple standard ELISA-test in which 15 kD or 32 kD protein or antigenic fragments thereof according to the invention are coated to the wall of the wells of an ELISA-plate. A method for the detection of such antibodies is e.g. incubation of 15 kD or 32 kD protein or antigenic fragments thereof with serum from mammals to be tested, followed by e.g. incubation with a labelled antibody against the relevant mammalian antibody. A colour reaction

30

can then reveal the presence or absence of antibodies against *Babesia canis* associated protein. Another example of a diagnostic test system is e.g. the incubation of a Western blot comprising the 15 kD or 32 kD protein or an antigenic fragment thereof according to the invention, with serum of mammals to be tested, followed by analysis of the blot.

Thus, another embodiment of the present invention relates to a diagnostic test for the detection of antibodies against *Babesia canis* associated antigenic material, characterised in that the test comprises a protein or an immunogenic fragment thereof according to the invention.

A diagnostic test based upon the detection of antigenic material of the specific 15 kD and 32 kD proteins amongst *Babesia canis* antigens and therefore suitable for the detection of *Babesia canis* infection can e.g. also be a standard sandwich-ELISA test.

In one example of such a test the walls of the wells of an ELISA plate are coated with antibodies directed against the 15 kD or 32 kD protein or immunogenic fragments thereof. After incubation with the material to be tested, labelled anti-*Babesia canis* antibodies are added to the wells. A colour reaction then reveals the presence of antigenic material from *Babesia canis*.

Therefore, still another embodiment of the present invention relates to a diagnostic test for the detection of *Babesia canis* associated antigenic material, characterised in that said test comprises antibodies against a protein or an immunogenic fragment thereof according to the invention.

The proteins or immunogenic fragments thereof according to the invention expressed as described above can be used to produce antibodies, which may be polyclonal, monospecific or monoclonal (or derivatives thereof). If polyclonal antibodies are desired, techniques for producing and processing polyclonal sera are well-known in the art (e.g. Mayer and Walter, eds. Immunochemical Methods in Cell and Molecular Biology, Academic Press, London, 1987).

Alternatively, antigenic material according to the invention can be used to produce monoclonal antibodies by immunising inbred mice by techniques also known in the art (Kohler and Milstein, Nature, 256, 495-497, 1975).

Still another embodiment of the invention relates to methods for the detection of antigenic material from *Babesia canis* in which the method comprises the incubation of serum, tissue or body fluids with antibodies against the 15 kD or the 32 kD protein or an antigenic fragment thereof according to the invention.

5

Examples

EXAMPLE 1: Molecular characterisation of *BCVIR* cDNA

5 Growth of *Babesia canis*

Isolates of *Babesia canis* (isolate A and B) were obtained from dogs naturally infected in France. They were maintained in *in vitro* culture according to Schetters *et al.* (1997, *Parasitology*, **115**, 485-493).

10 cDNA library construction, immunological screening and isolation of plasmid DNA.

A cDNA library from *Babesia canis* (isolate A) was constructed with the ZAP Express™ cDNA Gigapack® II Gold Cloning kit (Stratagene) as it was described for *B. rossi* in Carret *et al.* (1999, *Eur. J. Biochem.*, **265**, 1015-1021). The library was screened as described in Carret *et al.* (1999, *Eur. J. Biochem.*, **265**, 1015-1021) using the anti-GST60.1 (depleted of anti-*E. coli* antibodies), a polyclonal antibody directed against one member of the *Plasmodium falciparum* Pf60 multigene family (Carcy *et al.*, 1994, *Mol. Biochem. Parasitol.* **68**: 221-233), at a 1:100 dilution. Positive clones were visualised using goat anti-rabbit IgG conjugated to peroxidase (Sigma) at a 1:500 dilution, using 4-chloro-1-naphtol as chromogen (Sigma). They were purified and the pBK-CMV plasmids carrying the *BCVIR* cDNA were prepared as described in Carret *et al.* (1999, *Eur. J. Biochem.*, **265**, 1015-1021).

DNA sequencing

25 Nucleotid sequence was performed using the dideoxy chain termination method from alkali-denaturated double-strand templates according to Sanger *et al.* (1977, *Proc. Natl. Acad. Sci. USA* **74** (12):5463-5467) by Genome Express S.A (Zone Astec, Grenoble, France) on both strands of the selected plasmids using T3, T7 universal primers and oligonucleotides derived from established sequence of each strand.

Cloning of the 5' end of the *RCUTR* cDNA

obtained as follows. the cDNA library was used as a DNA template for a PCR experiment using the T3 sense universal primer derived from the vector sequence

(located 70 bp upstream of the *Eco*RI cloning site) and an antisense primer P2 (5'-ATGAGTCTATTGACTCCTTG-3') derived from the *BCVIR* cDNA sequence (nucleotide position 476-496 downstream of the *Eco*RI cloning site in the complete sequence of the cDNA, see Fig. 1 and Fig.2). The PCR was performed as described below (§ PCR amplification) using 1 µl of the cDNA library (10⁵ phages) as DNA template. The highest resulting PCR fragment (with a size of around 550 bp) was gel-extracted using gel extraction spin columns (Genomed), cloned in pGEM®-T vector using the pGEM®-T vector System II kit under the recommendations of Promega and sequenced on both strands as described behind, using T7 and SP6 universal primers (Genome Express).

PCR amplification

Amplifications were performed in a PTC-100™ Programmable Thermal Controller (MJ Research, INC.) as described in Carret *et al.* (1999, *J. Eukaryot. Microbiol.*, 46(3), 298-303) with the following conditions : 3 min. at 94°C, 30 cycles of 1 min. at 94°C, 1 min. at 55°C and 1 min. at 72°C, and then 5 min. at 72°C. Primers P1 (sense 5'-GACGTTTGATGTGATGAGGGAAGC-3'), P5 (sense 5'-AGGGAGCTGTCACGGAAGATT-3'), P2 (antisense) and P15.2 (antisense 5'-AATGACATACTCACAGGAAGC-3') were all derived from the *BCVIR* cDNA sequence (their respective position is indicated on the Fig.1) and commonly used for the molecular analysis of this sequence.

Genomic DNA analysis of the *BCVIR* cDNA sequence:

Genomic DNA extraction from *Babesia canis* cultures and Southern blot experiments were performed using standard procedures described in Maniatis/Sambrook (Sambrook J. *et al.*, Molecular cloning: a laboratory manual. ISBN 0-87969-309-6). The probes were obtained by performing PCR with the combinations of primers P1/P15.2 or P1/P2 using the pBK-CMV plasmid carrying the *BCVIR* cDNA as DNA template and they were labelled using Nick Translation kit under the recommendations of the manufacturer (Boehringer Mannheim). These probes were also tested on chromosomes of *B. canis* separated in 0.7% agarose (ICN) by pulse field gel electrophoresis (PFGE) and transferred onto Nylon membranes.

As control, hybridisation experiments were performed with a probe derived from the Bcc12D3 cDNA of *B. canis* A on the same membranes that were hybridized with probes derived from the *BCVIR* cDNA sequence. Dehybridisation was performed as recommended by Amersham Pharmacia Biotech.

5

RNA analysis of the *BCVIR* cDNA sequence:

Total RNA was extracted using the RNAgents^R Total RNA Isolation System according Promega and was used as starting material for RT/PCR and Northern experiments.

10

RT-PCR. RT-PCR was performed using the two-step protocol of the Enhanced Avian RT-PCR kit as described by the manufacturer (Sigma). Briefly, the first step (retrotranscription) was performed for 50 min. at 42°C using 250 pg of total RNA as template and 1 µM of specific cDNA primer (sense P1, sense P5 (5'-

15 AGGGAGCTGTCACGGAAGATT-3') or antisense P15.2), and the second step was performed at an annealing temperature of 50°C as described above (§PCR amplification), using one fifth of the first step reaction as a template. As controls, the first step was also performed without eAMV-RT for each primer tested for DNA contamination, and the experiments were also performed with specific primers (sense
20 Bcc12D3.1 and antisense Bcc12D3.2) derived from the Bcc12D3 cDNA of *B. canis* A.

Northern Blotting. For Northern blot experiments, total RNA was electrophoresed through gel containing formaldehyde and transferred to Nylon membrane as described
25 in Maniatis/Sambrook (Sambrook J. et al., Molecular cloning: a laboratory manual. ISBN 0-87969-309-6). Hybridisation was performed with a probe derived from a P1/P2 PCR fragment amplified on the *BCVIR* cDNA sequence and labelled with digoxigenin (DIG)-dUTP according the DIG High Prime DNA labelling kit (Boehringer Mannheim). Total RNA samples depleted in mRNA were also used for
30 hybridisation experiments and mRNA depletion was performed using PolyA Tract

Results:

Isolation and sequence of the *BCVIR* cDNA. Immunoscreening of a cDNA library of *B. canis* (isolate A), using the anti-GST60.1 antiserum directed against one

5 member of the Pf60 multigene family of *P. falciparum* (Carcy *et al.*, 1994, *Mol. Biochem. Parasitol.* **68**: 221-233) allowed to isolate and sequence the *BCVIR* cDNA.

The complete nucleotide sequence of *BCVIR* cDNA contained 1135 nucleotides (between *EcoRI* to *XhoI* cloning restriction sites) with a poly(A)₂₂ tail (Fig.1A and

Fig.2). An interesting feature of this cDNA is that two overlapping open reading
10 frames (ORF) are predicted (Fig.1A) : ORF1 between nucleotide 75 (ATG, initiation codon for Methionine) and 500 (TAA, stop codon) and ORF2 between nucleotides 448 (ATT codon for Isoleucine) and 933 (TGA, stop codon). These two ORF are overlapped by 52 nucleotides (position 448-500) and are +1 shifted (Fig.1A).

ORF1 encodes a predicted polypeptide of 141 residues (designated Bcvir15) starting
15 by a methionine with an expected molecular mass of 15.7 kDa and an iso-electric point of 10.9 (Fig. 2). ORF2 does not contain a consensus translation initiation sequence and started by Ile (Fig.1A). It encodes a predicted polypeptide of 133 residues with an expected molecular mass of 15.2 kDa. Due to the +1 shift between ORF1 and ORF2, these two ORFs present different amino acid sequences in the
20 overlapping region (Fig.1B). Sequence analysis of the overlap region of *BCVIR* cDNA suggested that such a +1 ribosomal frame shift in that region could produced an ORF1-ORF2 fusion protein of 285 residues (designated Bcvir32), resulting from the translation of a 859 nucleotides sequence (Fig.3). Bcvir32 would have predicted molecular mass of 32.3 kDa and an iso-electric point of 10.5.

25

DNA analysis: The presence of *BCVIR* cDNA sequence in the genome of *B. canis* (isolates A and B) was analysed by PCR and hybridisation experiments both on DNA (Southern blot) or chromosomes (PFGE) from these isolates.

30 **PCR experiments.** These were performed using 3 combinations of specific primers from the *BCVIR* cDNA sequence : P1/P2 and P5/P2 covering the ORF1 (Fig.4A and 4B, respectively), and P5/P15.2 covering the ORF2 and the overlapping region between these two ORF of the cDNA (Fig.4C). As controls, these combinations of primers allowed to amplify DNA fragment of around 360, 160 and 630 bp on the

cDNA *BCVIR* carried by the pBK-CMV plasmid (Fig.4A-C, lanes 5) and were negative on genomic DNA prepared from uninfected dogs (Fig.4A-C, lanes 4). These combinations of primers were unable to amplify genomic DNA fragments when they were tested on genomic DNA from the isolates A and B of *B. canis* (Fig.4, lanes 1 and 2, respectively) whereas, as control for the DNA preparations, specific primers derived from the Bcc12D3 cDNA of *B. canis* were able to specifically amplify the expected 1200 bp genomic DNA fragment on these preparations (Fig.4D, lane 1 and 2, respectively). Moreover, primers from the *BCVIR* cDNA sequence were negative in PCR on genomic DNA from the isolate F of *B. rossi* as were those derived from the Bcc12D3 cDNA (Fig.4, lanes 3).

Hybridisation experiments. hybridisation with probes synthesised using the PCR fragments P1/P2 and P1/P15.2 from the *BCVIR* cDNA sequence, were performed on genomic DNA of isolates A and B of *B. canis* and isolate F of *B. rossi*, either digested by various restriction enzymes (Southern blot) or on chromosomes (separated by PFGE). Whatever the hybridisation experiment performed, none of these probes allowed to detect the *BCVIR* sequence on genomic DNA of *B. canis*, whereas following de-hybridisation of these probes from the membranes, a probe derived from the Bcc12D3 cDNA sequence was able to detect its sequence on genomic DNA of *B. canis* (data not shown).

RNA analysis. Analysis of the *BCVIR* cDNA sequence was performed by RT-PCR (Fig.5, A-C) and hybridisation experiments (Northern blot) (Fig.6) using total RNA extracted from the isolate A of *B. canis*.

RT-PCR experiment. For the RT-PCR analysis of *BCVIR* sequence, analysis of the Bcc12D3 cDNA sequence was used as control (Fig.5D and 5E). As expected for this control, the 1200 bp DNA fragment from Bcc12D3 was amplified by PCR with the combination of primers Bcc12D3.1/Bcc12D3.2 when the first retrotranscription step was performed with antisense primer Bcc12D3.2 (Fig.5E, lane 5), but not when it was

performed with sense primer Bcc12D3.1 (Fig.5E, lane 4). For the *BCVIR* sequence, the expected 600 bp DNA fragment was amplified by PCR with the combination of primers P5/P15.2 PCR when the first step of retrotranscription was performed with the antisense primer P15.2 (Fig.5C, lane 3). However and

unexpectedly, this DNA fragment of 630 bp was also amplified by PCR with this P5/P15.2 combination of primers when the first step of retrotranscription was performed with the sense primer P5 (Fig.5B, lane 2). Identically, a unexpected 360 bp DNA fragment was amplified by PCR with the P1/P2 combination of primers when the first step of retrotranscription was performed with the sense primer P1 (Fig.5A, lane 1). The same experiments performed by omitting the e-AMV Retrotranscriptase enzyme in the retrotranscription steps did not allow to amplify DNA fragment by PCR with these combination of primers (Fig.5, lanes 1'-5'), demonstrating that the 360 and 630 bp DNA fragment amplified using the sense primers P1 or P5 in the retrotranscription step were not from genomic DNA origin, as previously observed (§ DNA analysis). Moreover, their sequencing and their hybridisation with a probe derived from the entire *BCVIR* cDNA sequence (Fig.5, II) certified their specificity.

Northern blot experiment. hybridisation performed onto total RNA extracted from *B. canis* (isolate A) with a probe corresponding to the ORF1 sequence of *BCVIR* cDNA allowed to detect three different RNA bands : a strong one of around 1400 bases length and two faint ones at around 1200 and 2800 bases length (Fig.6, lane 2). When hybridisation was performed with the same amount of starting total RNA but depleted in mRNA, the 1200 bases band disappeared (Fig.6, lane 3), indicating that this RNA band is a mRNA, in agreement with the size of the cDNA. It confirms also that the *BCVIR* cDNA sequence is carried by non messenger RNA (the 1400 and 2800 bases length bands), in agreement with the results obtained by RT-PCR when the retrotranscription step was performed with sense primer.

No signal was found when hybridised with the *BCVIR* cDNA sequence onto total RNA from *B. rossi* (Fig.6, lane 4) or the bovine species *B. divergens* (Fig.6, lane 1).

Conclusion: The molecular characterisation of the *BCVIR* cDNA sequence indicates that it is not from *B. canis* genomic DNA origin. It demonstrates that it is carried by a mRNA, but also by a non messenger RNA of 1.4 kb and 3 kb. It demonstrates also that the sequence from this cDNA is specifically found in the European isolates of *B. canis*, not in *Babesia rossi*.

Example 2: Biochemical characterisation of proteins encoded by the BCVIR cDNA sequence.

Expression and purification of GST-ORF1 and GST-ORF2 recombinant proteins in *E. coli*.

In order to express the products encoded by ORF1 and ORF2 from the *BCVIR* cDNA as recombinant proteins, and to produce specific antisera against them, the two ORFs were subcloned in the *EcoRI/XhoI* cut dephosphorylated pGEX-4T3 vector (Amersham Pharmacia Biotech) in frame with the glutathione-S-transferase (GST).

- 10 In order to produce Bcvir15 protein encoded by ORF1 (aa M¹-I¹⁴¹, Fig.2), the *EcoRI/XhoI* fragment containing the 1041 bp cDNA sequence (missing the 5' end of *BCVIR* cDNA) was excised from the pBK-CMV recombinant plasmid and subcloned into the vector. In order to obtain an antiserum specifically directed against the amino acid sequence deduced from the ORF2 sequence (aa sequence N¹⁵⁴-C²⁸⁵, Fig.3), *i.e.*
- 15 excluding the 52 bp overlapping region between the two ORF (nucleotide position 448-500 in Fig.1), an *EcoRI* restriction site was created in the ORF2 sequence by a PCR experiment. Thus, the 1041 bp *BCVIR* cDNA sequence carried by the pBK-CMV plasmid was used as a template DNA for the PCR experiment with the sense primer P_{Eco} (5'-ATGAGGAATTCGAACCGACTA-3' ; located at nucleotide position
- 20 529-549 on the complete *BCVIR* cDNA sequence, Fig.1 and Fig.2) and the antisense T7 universal primer derived from the vector sequence (located 75 bp downstream of the *XhoI* cloning site). The amplified fragment was cloned in pGEM[®]-T vector using the pGEM[®]-T vector System II kit under the recommendations of Promega. An *EcoRI/XhoI* fragment containing an around 600 bp DNA from ORF2 was released
- 25 from the recombinant plasmid and then subcloned into the vector. For both cloning, transformation with the ligation mixtures were performed into the *E. coli* BL21 cells and positive clones were selected using PCR with the specific primers from the *BCVIR* cDNA sequence.

- 30 Expression and purification of the recombinant GST-Bcvir15 protein and of a non recombinant GST from the pGEX-4T3 vector was performed according Smith and

Antibodies directed against the recombinant GST-Bcvir15 protein and GST-Bcvir25 protein were firstly isolated from total cell extract according the protocol of Nagai &

Thogersen (1987, In *Methods in enzymology* (ed. R. Wu and L. Grossman), 153, 461-481). They were then dissociated and loaded on a 12 % SDS-PAGE (Laemmli UK, 1974, *Nature (London)*, 287:680-685). Lastly, the recombinant GST-ORF2 was electro-eluted from the gel and was used to immunise a rabbit as described below.

5

Production of antibodies directed against GST-ORF1 and GST-ORF2 proteins.

Polyclonal antisera. Polyclonal antiserum raised against GST-ORF1 was produced both in mice and in a rabbit, and a polyclonal antiserum raised against GST-ORF2 was produced in a rabbit according E. Harlow & D. Lane (*Antibodies: a laboratory manual*. ISBN 2-907516-15-9). Rabbits (New Zealand White) were immunised with 50 µg of purified GST recombinant proteins emulsified in Freund's Complete Adjuvant (FCA) (Sigma) subcutaneously for the first injection, and by intramuscular injections for the two subsequent ones using Freund's Incomplete Adjuvant (FIA) (Sigma). Balb/C mice were intraperitoneally immunised with 15 µg of GST-ORF1 emulsified in FCA for the first injection and in FIA for the two subsequent ones. Injection of the animals was performed at 3-week intervals and the bleedings were done 8 days before each immunisation.

Production and screening of monoclonal antibodies. 48 h prior to fusion of splenocytes from the mice immunised with the recombinant GST-ORF1 with X-63 Ag8-653 myeloma cells, a last injection of 15 µg of GST-ORF1 was performed. The fusion process and cloning by limiting dilution for the isolation of monoclonal antibodies were performed according to Brown et al. (1979, *J. immunol. Methods*, 31:201-209). Hybridoma supernatants were screened by indirect immunofluorescence assay as described below.

Immunological methods:

Immunoblotting. Immunoblotting was performed with purified *B. canis* merozoites prepared as follows: parasitized erythrocytes from *in vitro* culture (20% of parasitaemia) were extensively washed with sterile RPMI 1640 medium (Life Technologies). The pellet was lysed with α-hemolysin from *S. aureus* (Sigma) at a final concentration of 36U/ml during 30 min. at 37°C. Free merozoites and remaining unlysed cells were pelleted by centrifugation (3500 g, 30 min.) and the supernatant

was discarded. The pellet was washed one time with TBS, resuspended in 2 ml of TBS and the merozoites were purified by centrifugation (3500 g, 30 min.) on a gradient of saccharose formed by a lower layer of 1.09 density and a upper layer of 1.02 density. Free merozoites were harvested at the interface, they were pelleted by centrifugation (15000 g, 10 min.) and washed 3 times with TBS. They were then processed for electrophoresis and the proteins were separated on a 15% SDS-PAGE. Immunoblotting was performed as described for *B. divergens* in Carcy *et al.* (1991, *Biol. Cell*, 72, 93-102) using a 1:100 dilution for the polyclonal antisera directed against GST-ORF1 and GST-ORF2 proteins.

Indirect immunofluorescence assays (IFA). IFA and double IFA were performed on *B. canis* parasitized erythrocytes (5% of parasitaemia) as described for *P. falciparum* in Grellier *et al.* (1994, *Biol. Cell*, 82, 129-138) using a 1:200 dilution for the polyclonal antisera directed against GST-ORF1 and GST-ORF2 proteins. For double IFA, the slides were first incubated with the anti-GST-ORF2 rabbit antiserum and then with mouse polyclonal or monoclonal antibodies directed against the GST-ORF1. Slides were mounted with Citifluor solution (Citifluor Ltd; London, UK) for limiting extinction fluorescence and the fluorescence was detected using a fluorescence microscope (Axioscope, Zeiss).

[³⁵S]-methionine radiolabelling of *B. canis* culture and immunoprecipitation.

[³⁵S]-methionine radiolabelling of *in vitro* cultures of *B. canis* with 50 µCi/ml (1200 Ci/mmol, Amersham-Pharmacia Biotech) and a 5% starting parasitaemia, and immunoprecipitation assays were performed as described for *B. divergens* in Carcy *et al.* (1991, *Biol. Cell*, 72, 93-102).

Protein phase separation by TX-114 treatment. The proteins of *B. canis* were phase separated in Triton X-114 (Sigma) as described for *B. divergens* in Precigout *et al.* (1991, *Infect. Immun.*, 59(8), 2799-2805) using 200 µl of [³⁵S]-methionine radiolabeled parasitized red blood cells (20% of parasitaemia) as starting material.

in vitro translation of the GST-ORF1 cDNA segment. To examine the possibility of a direct frame shifting was examined in an *in vitro*-translation system. *In vitro*-

translation products were synthesised using the TNT[®] Quick Coupled Transcription/Translation System (Promega) with a PCR DNA fragment containing the entire *BCVIR* cDNA sequence, a Kozacks sequence and a T7 promoting sequence at its 5' end for the *in vitro* translation. In order to amplify such a DNA fragment, the

5 PCR was performed on the pBK-CMV plasmid carrying the incomplete 1041 bp *BCVIR* cDNA as DNA template using the two following primers: a 90 mer sense oligonucleotide (primer P0 : 5'-

GGATCCTAATACGACTCACTATAGGGAGACCACCATGGAGTCGACATCAA
CAG ACCAACTTTGTTGCCGAGAACCGTCCCACCTTTGG-3'), containing a

10 Kozacks and a T7 promoting sequences but also the missing 5' end of the entire *BCVIR* cDNA sequence containing the starting ATG codon of the ORF1 (bolded in the primer sequence; nucleotide position 75-130 on the complete cDNA sequence; see Fig.1 and Fig.2), and the antisense primer P15.2 from the *BCVIR* cDNA sequence. The PCR was performed using 100 ng of the circular recombinant pBK-CMV plasmid

15 and the following conditions of amplification were used: 1 cycle of 3 min. at 94°C; 5 cycles of 1 min. at 94°C, 1 min. at 37°C and 1 min. at 72°C; 25 cycles of 1 min. at 94°C, 1 min. at 50°C and 1 min. at 72°C. The PCR products were gel extracted twice to ensure their purity and resuspended in nuclease free water at a concentration of 100 ng/μl. Then, they were added into a reticulocyte lysate to be translated using the

20 TNT[®] Quick Coupled Transcription/Translation System as described by the manufacturer (Promega). As control, the same experiment was performed by omitting the PCR fragment in the reaction. The total radiolabeled translated products (5 μl of the reaction) and immunoprecipitated products (from 20 μl of the reaction) with the anti-GST-ORF1 or the anti-GST-ORF2 or the anti-GST antisera and the pre-immune

25 rabbit serum were separated on a 15% SDS-PAGE.

Results:

30 **Purification of GST-ORF1 and GST-ORF2.** In order to produce polyclonal antibodies specifically directed against polypeptide encoded by ORF1 or ORF2, the 141 residues (¹M-¹⁴¹I) of ORF1 (encoded from ⁷⁵ATG to the stop codon TAA⁵⁰⁰) (Fig.2) and 132 residues (¹⁵⁴N-²⁸⁵C) of the ORF2 (encoded from nucleotide position

⁵³³ATT to the stop codon TGA⁹³³) (Fig.3) were cloned in the pGEX-4T3 vector in frame with the GST sequence. Overexpression of the GST-ORF1 and GST-ORF2 were performed in the BL21 *E. coli* strain (Fig. 7). Expression studies demonstrated that *E. coli* cultures containing the recombinant pGEX 4T3-ORF1 expressed, after
 5 IPTG induction, the GST-ORF1 protein as a soluble protein in the lysate of sonicated bacteria (Fig. 7A, lane 1). After elution, the soluble GST-ORF1 was purified and it presented the expected molecular mass of 40 kDa (Fig. 7A, lane 2). In contrast, the GST-ORF2 was insoluble in *E. coli* cultures containing the recombinant pGEX 4T3-ORF2 and the recombinant protein, with a molecular mass of 39 kDa, was mainly
 10 found in inclusion bodies (Fig. 7B, lane 5). In order to immunise a rabbit with the purified GST-ORF2, it was gel excised from an extract of purified inclusions bodies separated on 12% SDS/PAGE (Fig. 7B, lane 4).
 As control, a GST was purified after IPTG induction of an *E. coli* culture containing the native pGEX 4T3 (Fig. 7B, lane 3).

15 **Immunological studies:** Polyclonal antisera, raised against GST-ORF1 or GST-ORF2 or GST, were produced by immunising mice and rabbits in order to perform immunological studies of the cDNA encoded polypeptides. Two monoclonal antibodies directed against GST-ORF1 were also tested.

20 **Immunoprecipitation experiments.** in a first experiment, they were performed both on the total (Fig.8A) and exoantigens (Fig.8B) radiolabelled fractions of *B. canis* cultures (isolate A). In the total antigen fraction, the anti-GST-ORF1 strongly recognises a major product of 15 kDa (Bcvir15, indicated by an asterisk) and a minor
 25 product of around 32 kDa (putative Bcvir32, indicated by an arrow) (Fig.8A, lane 3). In the exoantigen fraction, only the minor product of around 32 kDa was detected faintly (Fig.8B, lane 3). The anti-GST-ORF2 was found not reactive against a 32 kD protein on the total or exoantigen fractions (Fig.8, lanes 4). Pre-immune rabbit and anti-GST antisera were not reactive on the antigens recognised with the anti-GST-
 30 ORF1 antiserum (Fig.8, lanes 1 and 2, respectively). In a second experiment, the

GST-ORF1 (Fig.9). It allowed to detect the 15 kDa protein in the hydrophilic fraction (Fig.9A, lane 2, indicated by an asterisk) and no reactivity was observed in the

hydrophobic fraction (Fig.9B, lane 2). As control, no reactivity was observed against the 15 kDa protein when the immunoprecipitation was performed with the pre-immune rabbit serum (Fig.9, lanes 1). MAb directed against GST-ORF1 were not reactive in immunoprecipitation experiments.

5

Western blot. Western blots were performed with purified merozoites from the isolate A of *B. canis*. The anti-GST-ORF1 specifically recognised the 15 kDa protein (Fig.10A, lane 2, indicated by an asterisk) but, in contrast to immunoprecipitation experiments, it did not recognise the minor 32 kDa product. However, the anti-GST-ORF2, which recognised different proteins in Western blot experiment, was able to detect such a 32 kDa antigen (Fig.10B, lane 2, indicated by an arrow). No protein of around 15 kDa (which could correspond to predicted 15.2 kDa polypeptide of the ORF2) was recognised by the anti-GST-ORF2. As controls, pre-immune rabbit serum from each antiserum were not reactive in western blot (Fig.10, lanes 1). MAb directed against GST-ORF1 were not reactive in Western blot.

Immunofluorescence assays. IFA tests were performed on fixed-blood smears from *in vitro* cultures of the isolates A of *B. canis* (Fig.11) with the anti-GST-ORF1 and the anti-GST-ORF2. The fluorescence obtained with the polyclonal rabbit anti-GST-ORF1 was associated with numerous intraparasitic vesicles, dispersed through all the parasite cytoplasm but excluding the parasite's nucleus (Fig.11, picture A). The same pattern of immunofluorescence was observed with the mAb directed against GST-ORF1. Moreover, a double IFA performed with the mAb directed against GST-ORF1 and the polyclonal rabbit anti-GST-ORF2 indicated that antigens recognised by anti-GST-ORF2 (Fig.11, picture B) co-localised with those recognised by the anti-GST-ORF1. As controls, anti-GST and pre-immune rabbit antiserum were not reactive in IFA tests. The same profile of fluorescence was observed when the IFA tests were performed on infected erythrocytes from the isolate B of *B. canis* (data not shown).

***In vitro* translation of the BCVIR cDNA.** An *in vitro* translation of the entire BCVIR cDNA sequence was performed in reticulocyte lysate in order to analyse, by immunoprecipitation experiments, its products of expression (Fig.12, lanes 3-4). Analysis of total expression products revealed that a major 15 kDa protein was expressed when the experiment was performed with the cDNA (Fig.12, lane 4;

indicated by an asterisk) but not when it was performed by omitting the cDNA as control (Fig.12, lane 1). This major 15 kDa protein was specifically immunoprecipitated by the anti-GST-ORF1 but also a minor protein of around 32 kDa (Fig.12, lane 3; indicated by an arrow). As controls, the pre-immune rabbit serum and the anti-GST antiserum were not reactive on these products (data not shown). Thus, it suggests that such a minor 32 kDa product could result from the predicted +1 frame shift mechanism. As was the case for immunoprecipitation assays previously described, the anti-GST-ORF2 was not reactive on this minor product (data not shown).

Conclusion: The anti-GST-ORF1 antiserum is able to detect two proteins, a hydrophilic major one of 15 kDa and a minor one of 32 kDa, the last one being detected in the supernatant of *in vitro* culture of *B. canis*. The anti-GST-ORF2 antiserum was also able to detect a 32 kDa protein, but was unable to detect a 15 kDa product. It indicates that the ORF1 of the *BCVIR* cDNA encodes for a 15 kDa protein (predicted 15.7 kDa product, designated Bcvir15) and that ORF2 is unable to synthesise the predicted 15.2 kDa product. Moreover, the detection of a 32 kDa protein commonly recognised by anti-GST-ORF1 and anti-GST-ORF2 antisera, suggest that it could correspond to the predicted 32.3 kDa product resulting from a +1 frame shift mechanism and allowing to produce a fused ORF1-ORF2 protein (designated Bcvir32). The *in vitro* translation of the *BCVIR* cDNA sequence indicated that such a mechanism might occur in the cDNA sequence. The low level of expression of a 32 kDa protein in comparison with the high expression level of Bcvir15 protein is in agreement with such a mechanism. Moreover, the co-localisation of proteins recognised by the two antisera in intraparasitic vesicles, indicating that the 32 kDa protein recognised by the two antisera is the same one, is consistent with such a frame-shift mechanism.

Example 3- *Babesia canis* growth-inhibition tests.

ADDITIONAL TESTS - INHIBITION

Before each experiment, the parasitaemia was adjusted to 1% with a 2% haematocrit (day D0). *B. canis* parasites (in conditions previously described in § growth of *B.*

canis) were *in vitro* cultured in 1 ml for experiments performed in 24 wells plates or in 2 ml for experiments performed in 6 wells plates. The ability of the anti-GST-ORF1 and anti-GST-ORF2 antisera to inhibit *B. canis* growth *in vitro* was determined by adding each antiserum to the medium at different concentrations (from 1 to 8%).

- 5 Controls were : i) incubation with the rabbit pre-immune serum or the rabbit anti-GST serum or with the irrelevant rabbit anti-GST-Bcc12D3 serum at same concentrations; ii) a *B. canis in vitro* culture without treatment.

The ability of each concentration of antiserum to inhibit *in vitro* growth of *B. canis* (isolate A and B), was performed in triplicate and determined by parasitaemia on

- 10 Giemsa-stained smears or by metabolic labelling with 50 μ Ci/ml of [3 H]-Hypoxanthine in aqueous solution (ICN), after increasing incubation times of 24 h (day D1) and 48 h (day D2) corresponding respectively to about 3 and 6 *B. canis* life cycles. After D1, half of the volume of the wells was collected and the culture medium was entirely removed and replaced by fresh medium supplemented with
- 15 antiserum (and [3 H]-Hypoxanthine for labelling experiment) at the same concentration used at D0. The collected infected erythrocytes (respectively 5 and 20 μ l in 24 and 6 wells plates) were pelleted by centrifugation at 1800 g for 5 min. and washed five times in TBS. For the determination of radioactivity incorporation, they were lysed in 500 μ l of RIPA buffer and placed 1 h on ice. The lysates were
- 20 centrifuged at 15000 g (15 min., 4 °C), 2 μ l of the supernatant were then added to 2 ml of Emulsifier-Safe (Packard, USA) and the incorporated radioactivity was measured.

In vitro inhibition assay with purified immunoglobulins.

- 25 Before each experiment, the parasitaemia was adjusted to 1% with a 2% haematocrit (day D0). *B. canis* and *B. rossi* parasites were cultured *in vitro* in 96 wells plates in 100 μ l of medium . Antisera used for inhibition assays (anti-GST-Bcvir15, anti-GST-Bcvir15 pre-immune, anti-GST-Bcc12D3, anti-GST-ORF2 antiserum and anti-GST (Respectively indicated as Bcvir15, T0, CONT, orf2 and GST in fig. 17)) were first
- 30 purified by FPLC on a HiTrap™ Protein G column (Amersham Pharmacia Biotech). Purified immunoglobulins were then dialyzed overnight at 4°C versus PBS 1X buffer and total immunoglobulins were titrated using the Coomassie Protein Reagent Kit (Pierce, USA). ELISA was used to evaluate the ratio of immunoglobulins directed either against a native GST protein or against each recombinant protein fused to GST

by comparison of reactivity of purified immunoglobulins on 0.1 µg of non recombinant GST or 0.1 µg of recombinant proteins coated in wells. All purified immunoglobulin samples were then adjusted at the same concentration of specific anti-GST immunoglobulins with RPMI. The ability of each sample to inhibit *in vitro* culture of *Babesia canis* or *Babesia rossi* was performed in triplicate with 1, 2, 4, 8 and 10 µg of specific immunoglobulins directed against the non GST part of the recombinant protein used to obtain the antisera. The growth inhibition was evaluated by metabolic labelling with 50 µCi/ml of [³H]-Hypoxanthine in aqueous solution (ICN), during 24 hours (about 3 cycles of *B. canis* life cycle). Then, the erythrocytes were blocked on a filter and washed using a Scatron apparatus. Afterward, the radioactivity associated with each well was read in 2 ml of Scintillant (Emulsifier Safe™ Packard, USA) by scintigraphy.

Reversion of the inhibition by recombinant GST-ORF1 protein

The ability of the recombinant GST-ORF1 to reverse the inhibitory effect of the anti-GST-ORF1 on growth of *B. canis* was tested on *in vitro* culture maintained as described above. Different concentrations of recombinant protein in the medium containing 8% of anti-GST-ORF1 were tested: 20 µg/ml and 40 µg/ml. Four controls were used: i) *B. canis in vitro* culture without treatment; ii) culture containing 20 µg/ml or 40 µg/ml of recombinant GST-ORF1; iii) a culture containing 8% of anti-GST-ORF1 antiserum.

The ability of each concentration of GST-ORF1 to reverse the inhibitory effect of the anti-GST-ORF1 antiserum was performed in triplicate and determined by parasitaemia on Giemsa-stained smears at D0, D1 and D2 as described above.

Results:

Inhibition test by antisera. In a first experiment, the ability of the different bleedings from the anti-GST-ORF1 antiserum (noted II, III and IV in the figure 13) to inhibit the *in vitro* growth of *B. canis* (isolate A) was tested by adding in the culture medium

the figure 13) compared to the control culture, where parasitaemia increased from 10% to 35% during the 2 days experiment (Fig.13E), a significant inhibitory effect of the anti-GST-ORF1 on the *in vitro* growth of *B. canis* was observed at D2 with the

second and the third bleedings of the antiserum at a concentration of 8% since the parasitaemia was of around 15% at D2 in these treated cultures (Fig.13D, III and IV, respectively). As control, no inhibitory effect of the pre-immune rabbit serum was observed whatever the concentration used (Fig.13, I).

- 5 Another experiment was performed in order to analyse the inhibitory effect of the anti-GST-ORF2 antiserum on the growth of *B. canis* (Fig.14). It was tested and compared to anti-GST-ORF1 antiserum, and pre-immune or anti-GST or irrelevant anti-GST-Bcc12D3 rabbit antisera were used as controls (Fig.14). All antisera were tested at the inhibitory concentration of anti-GST-ORF1 (8%). In that experiment, a
10 significant inhibitory effect of the anti-GST-ORF2 was also observed at D2 since the parasitaemia was of 22% in the treated culture whereas it was of 35% in the control culture. Moreover, the inhibitory effect of anti-GST-ORF2 was found lower than the one observed with the anti-GST-ORF1 (at D2 the parasitaemia is of around 15%). No inhibitory effect was observed by treating the culture with 8% of pre-immune or anti-
15 GST or anti-GST-Bcc12D3 rabbit antisera.

The 8% inhibitory effect of the anti-GST-ORF1 antiserum was also checked by measuring incorporation of [³H]-Hypoxanthine following metabolic labelling of treated and untreated cultures, and the experiment was performed on the isolates A and B of *B. canis* (Fig.15, A and B respectively). For both isolates and compared to
20 the control cultures, a significant inhibitory effect was observed at D2 since the radioactivity was of around 2000 cpm in the untreated cultures and respectively of 250 and 350 cpm in the isolate A and B. In contrast to parasitaemia counting which was unable to detect an inhibitory effect of the anti-GST-ORF1 at D1, this inhibitory effect was visible at this day by radioactivity counting. Indeed, radioactivity in the
25 control culture was of 1000 cpm whereas in treated cultures it was of 200 and 250 cpm respectively in the isolate A and B.

***In vitro* inhibition assay with purified immunoglobulins.**

The experiment described above was repeated with purified immunoglobulins, in
30 order to exclude non-immunoglobulin serum-components as a possible non-specific cause of growth inhibition. As is shown in figure 17, purified immunoglobulins are capable of inhibiting *in vitro* growth of both *Babesia canis* A and B.

Surprisingly, this implies that vaccines comprising one or both proteins according to the invention are capable of inhibiting parasitic growth of *B. canis* in general, regardless the serovar. As expected, *in vitro* growth of *B. rossi* M is not inhibited.

- 5 **Reversion of the inhibition.** In order to confirm the inhibitory effect of anti-GST-ORF1 on the *in vitro* growth of *B. canis*, an experiment was conducted to reverse this effect by adding in the culture different concentration of purified recombinant GST-ORF1 (Fig.16). The two concentrations used were 20 and 40 µg/ml of GST-ORF1. This experiment showed that addition of GST-ORF1 protein in the medium of a
- 10 culture treated with 8% of anti-GST-ORF1 allowed a 50 and 100% reversion of the inhibition process respectively at concentration of 20 and 40 µg/ml (Fig.16, C and D respectively). Such a concentration of GST-ORF1 added in a non treated culture had no effect on the growth of the parasite (Fig.16, A and B respectively). Untreated culture and a treated culture with 8% of anti-GST-ORF1 were used as controls for this
- 15 experiment (Fig.16, E and F, respectively).

- Conclusion:** Both anti-GST-ORF1 and anti-GST-ORF2, at a concentration of 8% were able to inhibit, significant and specifically, the *in vitro* growth of *B. canis* isolate. This effect was already observed at D1. A concentration of 40 µg/ml of
- 20 recombinant GST-ORF1 was able to totally reverse this inhibitory effect. Moreover, filtration of anti-GST-ORF1 and anti-GST antisera indicated that 8% of each antiserum represents a respective concentration of immunoglobulins of 100 and 50 µg/ml of culture (data not shown). An ELISA test, performed on coated recombinant GST, indicated that half of the immunoglobulins from the anti-GST-ORF1 were
- 25 directed against the GST part of the recombinant GST-ORF1 (data not shown), it means that the inhibitory effect observed at 8% of anti-GST-ORF1 on the *in vitro* growth of *B. canis* was obtained by 50 µg/ml of immunoglobulins specifically directed against the protein BcVir15. At least, this is consistent with the 40 µg of recombinant GST-ORF1 necessary to reverse the inhibitory effect of the antiserum
- 30 anti-GST-ORF1.

for which the genetic information is present in *B. canis* but not encoded by the *B. canis* genomic DNA play an important role in the virulence of the Apicomplexan

parasite *B. canis*. Vaccines based upon these proteins or nucleic acid sequences encoding these proteins are shown to play an important role in the inhibition of *B. canis* parasite growth, regardless the serovar.

Figure 1: Schematic representation of the *BCVIR* cDNA

A: Schematic representation of entire *BCVIR* cDNA with predicted ORFs represented by boxes and sequence derived primers represented with arrows

- 5 B: Representation of the predicted overlapping sequence and its deduced amino acid sequence

Figure 2: Nucleic sequence of the complete *BCVIR* cDNA and deduced amino acid sequence of the Bcvir15.

- 10 The common amino acid sequence between Bcvir15 and the putative Bcvir32 is represented in regular upper case. A GRAM-positive anchoring hexapeptide is overlined. The overlapping sequence between the two predicted ORF of *BCVIR* cDNA are indicated in bold italic. Cloning restriction sites are in bold case.

- 15 **Figure 3: Nucleic sequence of the complete *BCVIR* cDNA and deduced amino acid sequences of the putative Bcvir32 protein.**

The overlapping region between ORF1 and ORF2 of the *BCVIR* cDNA is indicated in **bold italic**. Cloning restriction sites are in bold case.

- 20 **Figure 4: PCR analysis of *BCVIR* cDNA sequence on genomic DNA of *B. canis***

Genomic DNA analysed were extracted from isolate A of *B. canis* (lane 1), isolate B of *B. canis* (lane 2), isolate F of *B. rossi* (lane 3) and uninfected dog blood sample (lane 4). The pBK-CMV plasmid carrying the *BCVIR* cDNA sequence was used as positive control (lane 5). PCR were performed using the combination of primers

- 25 P1/P2 (A), P5/P2 (B), P5/P15.2 (C) and Bcc12D3.1/Bcc12D3.2 (D)

Figure 5: RT-PCR analysis of the *BCVIR* sequence

The experiment was performed on total RNA isolated from *B. canis* (isolate A). P1 (lane 1), P5 (lane 2), P15.2 (lane 3); all derived from *BCVIR* sequence and Bcc12D3.1 (lane 4) or Bcc12D3.2 (lane 5) of *Bcc12D3* sequence were used as initial primers for

the first step of amplification. The first step of amplification was performed with the

combination of primers P1/P2 (A), P5/P2 (B), P5/P15.2 (C) and Bcc12D3.1/Bcc12D3.2 (D).

5') The second step of amplification was performed by PCR with the combination of primers P1/P2 (A), P5/P15.2 (B, C) and Bcc12D3.1/Bcc12D3.2 (D). Amplified

products were separated on 0.8% agarose gel and visualised by Bet staining under UV light (I) or they were transferred on to Nitrocellulose membrane and hybridised with a total *BCVIR* cDNA sequence used as a probe in order to check the specificity of the amplified PCR products (II).

5

Figure 6: Northern blot analysis of the *BCVIR* cDNA sequence

Hybridisation was performed using the PCR derived P1/P2 probe of *BCVIR* cDNA sequence on total RNA from the isolate Rouen 1987 of *B. divergens* (lane 1), total RNA from the isolate A of *B. canis* (lane 2), non messenger RNA from the isolate A of *B. canis* (lane 3) and total RNA the isolate F of *B. rossi* (lane 4).

10

Figure 7: Over expression and purification of the GST-ORF1 and GST-ORF2 recombinant proteins from *E. coli*

Proteins were purified from BL21 (DE3) cells after 3 h IPTG induction as described, separated on a SDS/12% polyacrylamide gel, and stained with Coomassie brilliant blue.

15

Lanes : 1A, *E. coli* expressing GST-ORF1 lysate ; 1B, purified GST-ORF1; 3B control purified native GST; 4B purified GST-ORF2; inclusion bodies lysate from *E. coli* expressing GST-ORF2.

20

Figure 8: Immunoprecipitation of [³⁵S] methionine labelled *Babesia canis* isolate A antigens

Radiolabelled Total antigens (A) or Exoantigens (B) were immunoprecipitated respectively by rabbit pre-immune sera (1), anti-GST rabbit polyclonal serum (2), anti-GST-ORF1 rabbit polyclonal serum (3) and anti-GST-ORF2 rabbit polyclonal serum. Arrows and asterisk indicate respectively the location of the BcVir15 and the putative BcVir32.

25

Figure 9: Immunoprecipitation of [³⁵S] methionine labelled *Babesia canis* A strain TX-114 partitioned antigens

30

The [³⁵S] methionine labelled *in vitro* cultures of *Babesia canis* A isolate were solubilized in TX-114, and after a phase separation, the hydrophilic were recovered in the aqueous phase and the amphiphilic antigens were recovered in the detergent phase. Pre-immune rabbit serum (1) or anti-GST-ORF1 rabbit polyclonal serum (2)

were used to immunoprecipitated either the aqueous-phase proteins (A) or the detergent phase proteins (B). Arrow indicate the location of the BcVir15.

Figure 10: Immunoblot analysis of SDS-PAGE separated *Babesia canis*

5 merozoite antigens

The experiment was performed on purified merozoites of the isolate A of *Babesia canis*. Antigens were separated on a 12% SDS/polyacrylamide gel and transferred onto nitrocellulose sheets. Incubation was done at a 1 : 100 dilution with pre-immune rabbit serum (1A and 1B), with anti-GST-ORF1 rabbit polyclonal serum (2A) or with
 10 anti-GST-ORF2 rabbit polyclonal serum (2B). The position of the BcVir15 and BcVir32 are indicated respectively by an asterisk and an arrow.

Figure 11: Localisation of *BCVIR* sequence expressed products.

IFA tests were performed with the isolate A of *Babesia canis*.

- 15 (A) fluorescence associated with anti-GST-ORF1
 (B) fluorescence associated with anti-GST-ORF2

Figure 12: Analysis of *in vitro* translated products.

Radiolabelled *In vitro* products translated from P0/P15.2 PCR amplification on pBK-CMV *BCVIR* cDNA were separated on a 12% SDS-PAGE. Total products (4) or
 20 products immunoprecipitated by anti-GST-ORF1 (3) were analysed. Total expression product of reticulocyte lysate without DNA (1) or immunoprecipitated product from a reticulocyte lysate without DNA (2) were used as controls. Asterisk localise BcVir15 and arrow localise BcVir32.

25

Figure 13: evaluation of the capacity of different concentration of anti-GST-ORF1 rabbit polyclonal sera to inhibit *Babesia canis* A strain *in vitro* culture.

Different bleeding (respectively first (II), second (III) and third (IV) anti-GST-ORF1 rabbit polyclonal serum were added at a final concentration of 1% (A), 2% (B), 4%
 30 (C) and 8% (D) to *Babesia canis* A *in vitro* culture. Pre-immune serum (I) was used as

Figure 14: Inhibition of *Babesia canis* A strain *In vitro* culture by different rabbit polyclonal sera

Sera were tested for their ability to inhibit parasite growth when added at a rate of 8% on *Babesia canis* A strain *in vitro* culture. D0, D1, D2 indicate respectively parasitaemia values at the start of the test, after 24 hours of culture growth and after 48 hours of culture growth. Tested sera were rabbit polyclonal anti-GST-ORF1 and anti-GST-ORF2 and control sera were rabbit pre-immune serum, rabbit polyclonal anti-GST and rabbit polyclonal anti-GST-Bcc12D3. Control lane indicate culture in standard conditions without addition of any rabbit serum.

Figure 15: Inhibition test of *Babesia canis in vitro* culture evaluated by [³H] hypoxanthine metabolic labelling

Anti-GST-ORF1 and control pre-immune rabbit sera were all tested at 8% final concentration on *Babesia canis* A strain (I) and *Babesia canis* B strain (II). Control lane indicate in both case parasites grown in standard conditions.

Figure 16: Inhibition test against GST-ORF1 recombinant protein

The recombinant GST-ORF1 protein was tested for its ability to revert the capacity of rabbit polyclonal anti-GST-ORF1 serum to inhibit *Babesia canis* A strain growth *in vitro*. The protein was tested at 20µg/ml (C) or 40µg/ml (D) against 8 % of anti-GST-ORF1 rabbit polyclonal serum. Controls were protein alone added in the culture at a rate of 20µg/ml (A) or 40µg/ml (B), culture in standard conditions (E) and culture with 8% of anti-GST-ORF1 rabbit polyclonal serum (F).

Figure 17: Inhibition test of *Babesia canis in vitro* culture with purified immunoglobulins evaluated by [³H] hypoxanthine metabolic labelling.

The ability of purified immunoglobulins (anti-GST-Bcvir15, anti-GST-Bcvir15 pre-immune, anti-GST-Bcc12D3, anti-GST-ORF2 antiserum and anti-GST (Respectively indicated as Bcvir15, T0, CONT, orf2 and GST in fig. 17) to inhibit *in vitro* culture of *Babesia canis* or *Babesia rossi* was performed in triplicate with 1, 2, 4, 8 and 10 µg of specific immunoglobulins. Growth inhibition was evaluated by metabolic labelling with 50 µCi/ml of [³H]-Hypoxanthine.

Claims

06. 03. 2001

(38)

- 1) Nucleic acid sequence encoding a 15 kD *Babesia canis* associated protein or an immunogenic fragment of said protein, said protein or immunogenic fragment thereof having at least 80 %, preferably 90 %, more preferably 95 % homology with the amino acid sequence as depicted in SEQ ID NO: 2.
 - 2) Nucleic acid sequence encoding a 32 kD *Babesia canis* associated protein or an immunogenic fragment of said protein, said protein or immunogenic fragment thereof having at least 80 %, preferably 90 %, more preferably 95 % homology with the amino acid sequence as depicted in SEQ ID NO: 4.
 - 3) cDNA fragment comprising a nucleic acid sequence according to claim 1 or 2.
 - 4) Recombinant DNA molecule comprising a nucleic acid sequence according to claim 1 or 2 or a cDNA fragment according to claim 3, under the control of a functionally linked promoter.
 - 5) Live recombinant carrier comprising a cDNA fragment according to claim 3 or a recombinant DNA molecule according to claim 4.
 - 6) Host cell comprising a nucleic acid sequence according to claim 1 or 2, a cDNA fragment according to claim 3, a recombinant DNA molecule according to claim 4 or a live recombinant carrier according to claim 5.
 - 7) *Babesia canis* associated protein, said protein having a molecular weight of 15 kD and comprising an amino acid sequence that is at least 80 % homologous to the amino acid sequence as depicted in SEQ ID NO: 2 or an immunogenic fragment of said protein.
- amino acid sequence as depicted in SEQ ID NO: 2, or an immunogenic fragment of

said protein.

9) *Babesia canis* associated protein, said protein having a molecular weight of 32 kD and comprising an amino acid sequence that is at least 80 % homologous to the amino acid sequence as depicted in SEQ ID NO: 4 or an immunogenic fragment of said protein.

10) *Babesia canis* associated protein according to claim 9, wherein the amino acid sequence is at least 85 %, preferably 90 %, more preferably 95 % homologous to the amino acid sequence as depicted in SEQ ID NO: 4, or an immunogenic fragment of said protein.

11) *Babesia canis* associated protein according to claims 7-10 for use in a vaccine.

12) Use of a *Babesia canis* associated protein according to claims 7-10 for the manufacturing of a vaccine for combating *Babesia canis* infections.

13) Vaccine for combating *Babesia canis* infections, characterised in that it comprises a nucleic acid sequence according to claim 1 or 2, a cDNA fragment according to claim 3, a recombinant DNA molecule according to claim 4, a live recombinant carrier according to claim 5, a host cell according to claim 6 or a protein according to claims 7-10, and a pharmaceutically acceptable carrier.

14) Vaccine according to claim 13, characterised in that it comprises an adjuvant.

15) Vaccine according to claim 13 or 14, characterised in that it comprises an additional antigen derived from a virus or micro-organism pathogenic to dogs or genetic information encoding said antigen.

16) Vaccine according to claim 15, characterised in that said virus or micro-organism pathogenic to dogs is selected from the group of *Ehrlichia canis*, *Babesia gibsoni*, *vogeli*, *rossi*, *Leishmania donovani*-complex, Canine parvovirus, Canine distempervirus, *Leptospira interrogans serovar canicola*, *icterohaemorrhagiae*,

pomona, *grippotyphosa*, *bratislava*, Canine hepatitisvirus, Canine parainfluenzavirus, rabies virus, *Hepatozoon canis* and *Borrelia burgdorferi*

17) Vaccine for combating *Babesia canis* infections, characterised in that it comprises antibodies against a protein according to claims 7-10, or an immunogenic fragment thereof.

18) Method for the preparation of a vaccine according to claims 13-16, said method comprising the admixing of a nucleic acid sequence according to claim 1 or 2, a cDNA fragment according to claim 3, a recombinant DNA molecule according to claim 4, a live recombinant carrier according to claim 5, a host cell according to claim 6 or a protein according to claims 7-10 and a pharmaceutically acceptable carrier.

19) Method for the preparation of a vaccine according to claim 17, said method comprising the admixing of said antibodies and a pharmaceutically acceptable carrier.

20) Diagnostic test for the detection of *Babesia canis* associated RNA characterised in that the test comprises a nucleic acid sequence that is at least 70 % homologous to the nucleic acid sequence as depicted in SEQ ID NO: 1 or 3 or a nucleotide sequence that is complementary to said nucleic acid sequence, or a fragment thereof having a length of at least 12, preferably 15, more preferably 18 nucleotides.

21) Diagnostic test for the detection of antibodies against *Babesia canis* associated antigenic material, characterised in that said test comprises a protein or an immunogenic fragment thereof as defined in claims 7-10.

22) Diagnostic test for the detection of *Babesia canis* associated antigenic material, characterised in that said test comprises antibodies against a protein or an immunogenic fragment thereof as defined in claims 7-10.

1. The first part of the document discusses the importance of maintaining accurate records of all transactions and activities. It emphasizes that proper record-keeping is essential for transparency and accountability, particularly in financial matters. The text suggests that organizations should implement robust systems to track every detail, from budget allocations to expenditure reports.

2. The second section addresses the challenges faced by organizations in managing their resources effectively. It highlights the need for strategic planning and the allocation of funds based on long-term goals. The author argues that without a clear vision and a structured approach, organizations risk mismanaging their assets and failing to achieve their intended outcomes.

3. The third part of the document focuses on the role of leadership in driving organizational success. It stresses that leaders must be proactive in identifying opportunities and addressing challenges. The text provides insights into how effective leaders can inspire their teams, foster innovation, and ensure that the organization remains agile in the face of changing circumstances.

4. The fourth section explores the importance of communication in building a strong organizational culture. It notes that clear and consistent communication is vital for ensuring that all employees are aligned with the organization's mission and values. The author suggests that leaders should encourage open dialogue and provide regular updates to keep the team informed and motivated.

5. The final part of the document discusses the importance of continuous improvement and learning. It encourages organizations to regularly evaluate their performance and seek ways to enhance their processes. The text suggests that by embracing a growth mindset and being open to feedback, organizations can achieve sustained success and adapt to the ever-changing business environment.

Abstract

06. 03. 2001

The present invention relates to nucleic acid sequences encoding novel *Babesia canis* associated proteins and to cDNA fragments, recombinant DNA molecules and live
5 recombinant carriers comprising these sequences. Furthermore, the invention relates to host cells comprising such nucleic acid sequences, cDNA fragments, recombinant DNA molecules and live recombinant carriers. Also, the invention relates to proteins encoded by these nucleotide sequences, to vaccines for combating *Babesia canis* infections comprising these proteins or genetic material encoding these proteins and
10 methods for the preparation of vaccines. Another embodiment of the invention relates to these *Babesia canis* associated proteins for use in vaccines and to the use of the *Babesia canis* associated proteins in the manufacture of vaccines. Finally the invention relates to diagnostic tools for the detection of *Babesia canis* associated nucleic acid sequences, for the detection of *Babesia canis* associated antigens and for
15 the detection of antibodies against *Babesia canis* associated antigenic material.

the 1990s, the number of people in the UK who are aged 65 and over has increased from 10.5 million to 12.5 million, and the number of people aged 75 and over has increased from 4.5 million to 6.5 million (Office for National Statistics 2000). The number of people aged 65 and over is projected to increase to 15.5 million by 2020, and the number of people aged 75 and over to 8.5 million (Office for National Statistics 2000). The increase in the number of people aged 65 and over is expected to be due to a combination of factors, including a decline in the birth rate, a decline in the death rate, and a decline in the rate of emigration.


The increase in the number of people aged 65 and over is expected to have a significant impact on the UK's health and social care system. The number of people aged 65 and over who are in need of health and social care services is expected to increase significantly in the coming years. This is due to a number of factors, including a decline in the birth rate, a decline in the death rate, and a decline in the rate of emigration. The increase in the number of people aged 65 and over is expected to have a significant impact on the UK's health and social care system.

The increase in the number of people aged 65 and over is expected to have a significant impact on the UK's health and social care system. The number of people aged 65 and over who are in need of health and social care services is expected to increase significantly in the coming years. This is due to a number of factors, including a decline in the birth rate, a decline in the death rate, and a decline in the rate of emigration. The increase in the number of people aged 65 and over is expected to have a significant impact on the UK's health and social care system.

The increase in the number of people aged 65 and over is expected to have a significant impact on the UK's health and social care system. The number of people aged 65 and over who are in need of health and social care services is expected to increase significantly in the coming years. This is due to a number of factors, including a decline in the birth rate, a decline in the death rate, and a decline in the rate of emigration. The increase in the number of people aged 65 and over is expected to have a significant impact on the UK's health and social care system.

The increase in the number of people aged 65 and over is expected to have a significant impact on the UK's health and social care system. The number of people aged 65 and over who are in need of health and social care services is expected to increase significantly in the coming years. This is due to a number of factors, including a decline in the birth rate, a decline in the death rate, and a decline in the rate of emigration. The increase in the number of people aged 65 and over is expected to have a significant impact on the UK's health and social care system.

The increase in the number of people aged 65 and over is expected to have a significant impact on the UK's health and social care system. The number of people aged 65 and over who are in need of health and social care services is expected to increase significantly in the coming years. This is due to a number of factors, including a decline in the birth rate, a decline in the death rate, and a decline in the rate of emigration. The increase in the number of people aged 65 and over is expected to have a significant impact on the UK's health and social care system.



ORF2 I F V T L D L L R A K E S I D S S K K A L R

5' GCAGAAATTTTCGTCACCTTAGATCTCTTAAGAGCCAAAGGAGTCAATAGACTCATCTAAAGAGGCACTACGTA 3'

ORF1 A E Y F R H L R S L K S Q G V N R L I stop

EPO - DG 1
06. 03. 2001

(38)

Figure 1

EcoRI
gaattcggcacgagccctgctatactgtgctttgcaactaactccatcgtaataattttaatataataataaa 72

M E S T S T T T N F V A E N R P T 17
 gg ATG GAG TCG ACA TCA ACA ACG ACC AAC TTT GTT GCC GAG AAC CGT CCC ACC 125

F G E T F D V M R E A L L R V K S S 35
 TTT GGT GAG ACG TTT GAT GTG ATG AGG GAA GCT TTG CTT CGT GTA AAG TCC TCT 179

E R L A M L R A L A G M C G H R V L 53
 GAA CGC TTG GCA ATG CTC AGA GCG CTT GCA GGA ATG TGC GGT CAC CGC GTC CTT 233

P G T G A S A I A A T V T P K G A S 71
 CCT GGC ACT GGT GCT TCT GCG ATA GCG GCA ACG GTA ACC CCA AAG GGG GCT TCG 287

M K L K P P R P Q S T K S P E L R E 89
 ATG AAG CTT AAA CCA CCG CGT CCG CAG TCA ACG AAG TCT CCG GAG CTC AGG GAG 341

L S R K I R E M N K T I S Q E S A R 107
 CTG TCA CGG AAG ATT CGC GAA ATG AAT AAG ACT ATA AGT CAG GAA TCA GCT CGG 395

V N H R L P E G H P L L E K R A E Y 125
 GTA AAC CAC CGG TTG CCG GAA GGC CAC CCT CTC TTA GAG AAG CGG GCA GAA **TAT** 449

F R H L R S L K S Q G V N R L I * 141
TTT CGT CAC CTT AGA TCT CTT AAG AGC CAA GGA GTC AAT AGA CTC ATC TAA G 501
 aaggcactacgtaggtaccgtgcctctatgaggaatacgaaccgactagtcacaaatagacgaccagttcta 573
 ccaaaggtagagcctgactctaattctaccattcggccagcgacggagtcgcatgacaacgtggaatcttaga 645
 ccacgccggacgggttatccgtcaaatggtactttggcagttacggaactcctgatctcgatttatagatca 717
 aacttctacacctgaagggtggtcgaggaaggagatgtacgtgctgcaacacccataaggagcaagcttg 789
 ctactcctatccggttacctccagctatatcggtgcaactgcactcagttggaaggctctgtattcgtagaatac 861
 tgcaaaaccaggatagcggtcgaggcacgcctcaccggactacgtccgaggggtgaccctaacgggctgctga 933
 actaggttcagccagcgcttcctgtgagtatgtcattccgggtccttcggggcccgggccagtttcgactgg 1005
 tgtagggtttgccctactagagtactttgcgacgccgaagcgctccggttcaaaagaacgcgcaagccctagca 1077
 gagaatgcgagggcatgactcttcgagtcacaaaaaaaaaaaaaaaaaaaaa**ctcgag** 1135

XhoI

Figure 2

EcoRI

gaattcggcacgagccctgctatactgtgctttgcaactaactccatcgtaataatttaataataataaaa72

M	E	S	T	S	T	T	T	N	F	V	A	E	N	R	P	T	17
gg	ATG	GAG	TCG	ACA	TCA	ACA	ACG	ACC	AAC	TTT	GTT	GCC	GAG	AAC	CGT	CCC	ACC 125
F	G	E	T	F	D	V	M	R	E	A	L	L	R	V	K	S	S 35
TTT	GGT	GAG	ACG	TTT	GAT	GTG	ATG	AGG	GAA	GCT	TTG	CTT	CGT	GTA	AAG	TCC	TCT 179
E	R	L	A	M	L	R	A	L	A	G	M	C	G	H	R	V	L 53
GAA	CGC	TTG	GCA	ATG	CTC	AGA	GCG	CTT	GCA	GGA	ATG	TGC	GGT	CAC	CGC	GTC	CTT 233
P	G	T	G	A	S	A	I	A	A	T	V	T	P	K	G	A	S 71
CCT	GGC	ACT	GGT	GCT	TCT	GCG	ATA	GCG	GCA	ACG	GTA	ACC	CCA	AAG	GGG	GCT	TCG 287
M	K	L	K	P	P	R	P	Q	S	T	K	S	P	E	L	R	E 89
ATG	AAG	CTT	AAA	CCA	CCG	CGT	CCG	CAG	TCA	ACG	AAG	TCT	CCG	GAG	CTC	AGG	GAG 341
L	S	R	K	I	R	E	M	N	K	T	I	S	Q	E	S	A	R 107
CTG	TCA	CGG	AAG	ATT	CGC	GAA	ATG	AAT	AAG	ACT	ATA	AGT	CAG	GAA	TCA	GCT	CGG 395
V	N	H	R	L	P	E	G	H	P	L	L	E	K	R	A	E	Y 125
GTA	AAC	CAC	CGG	TTG	CCG	GAA	GGC	CAC	CCT	CTC	TTA	GAG	AAG	CGG	GCA	GAA	TAT 449
F	V	T	L	D	L	L	R	A	K	E	S	I	D	S	S	K	K 143
(T)	TTC	GTC	ACC	TTA	GAT	CTC	TTA	AGA	GCC	AAG	GAG	TCA	ATA	GAC	TCA	TCT	AAG 504
A	L	R	R	Y	R	A	S	M	R	N	T	N	R	L	V	H	N 161
GCA	CTA	CGT	AGG	TAC	CGT	GCC	TCT	ATG	AGG	AAT	ACG	AAC	CGA	CTA	GTG	CAC	AAT 558
R	R	P	V	L	P	K	V	E	P	D	S	N	L	P	F	G	Q 179
AGA	CGA	CCA	GTT	CTA	CCA	AAG	GTA	GAG	CCT	GAC	TCT	AAT	CTA	CCA	TTC	GGC	CAG 612
R	R	S	R	M	T	T	W	N	L	R	P	R	R	T	G	Y	P 197
CGA	CGG	AGT	CGC	ATG	ACA	ACG	TGG	AAT	CTT	AGA	CCA	CGC	CGG	ACG	GGT	TAT	CCG 666
S	N	G	T	L	A	V	T	E	L	L	I	S	I	Y	R	S	N 215
TCA	AAT	GGT	ACT	TTG	GCA	GTT	ACG	GAA	CTC	CTG	ATC	TCG	ATT	TAT	AGA	TCA	AAC 720
F	Y	T	L	K	V	V	E	E	G	R	C	T	C	C	N	T	H 233
TTC	TAC	ACC	TTG	AAG	GTG	GTC	GAG	GAA	GGG	AGA	TGT	ACG	TGC	TGC	AAC	ACC	CAT 774
K	E	Q	A	L	L	L	L	S	G	Y	L	Q	L	Y	R	A	L 251
AAG	GAG	CAA	GCT	TTG	CTA	CTC	CTA	TCC	GGT	TAC	CTC	CAG	CTA	TAT	CGT	GCA	CTG 828
H	S	V	G	R	S	V	F	V	E	Y	C	K	T	R	I	C	V 269
CAC	TCA	GTT	GGA	AGG	TCT	GTA	TTC	GTA	GAA	TAC	TGC	AAA	ACC	AGG	ATA	TGC	GTC 882
E	A	R	L	T	G	L	R	P	R	V	T	L	T	G	C	*	285
GAG	GCA	CGC	CTC	ACC	GGA	CTA	CGT	CCG	AGG	GTG	ACC	CTA	ACG	GGC	TGC	TGA	A 934
ctaggttcagccagcgccttcctgtgagtatgtcattccgggtccttcggggcccgccagtttcgactggt	1006																
qtaggtttgccctactagagtacttgcgaacgcgaagcgccttcggttcaaaagaacgcgcgaagccctagcag	1078																
agaaatgcgagggcatgactcttcgagtcacaaaaaaaaaaaaaaaaaaaaactcgag	1135																

XhoI

Figure 3




Figure 4




Figure 5




Figure 6




Figure 7




Figure 8




Figure 9.




Figure 10




Figure 11




Figure 12





Figure 13






Figure 16






Figure 15


Inhibition *in vitro* *B. canis* A



Inhibition *in vitro* *B. canis* B



Inhibition *in vitro* *B. rossi* M



1. The first part of the document discusses the importance of maintaining accurate records of all transactions and activities. It emphasizes that proper record-keeping is essential for transparency and accountability, particularly in financial matters. The text outlines various methods for organizing and storing data, including digital databases and physical filing systems.

2. The second section focuses on the role of communication in project management. It highlights the need for clear, concise, and timely communication between all stakeholders involved in a project. The author provides several strategies for effective communication, such as regular meetings, status reports, and the use of collaborative tools.

3. The third part of the document addresses the challenges of resource allocation and management. It discusses how to identify and prioritize tasks, allocate resources efficiently, and monitor progress. The text also touches upon the importance of flexibility and adaptability in the face of changing circumstances.

4. The final section discusses the importance of risk management. It explains how to identify potential risks, assess their impact, and develop mitigation strategies. The author stresses that proactive risk management is crucial for ensuring the success of any project or initiative.

SEQUENCE LISTING

EPO - DG 1

06. 03. 2001

(38)

<110> AKZO Nobel N.V.

<120> Babesia canis vaccine

<130> Bccvirvaccine

<140>

<141>

<160> 10

<170> PatentIn Ver. 2.1

<210> 1

<211> 1135

<212> DNA

<213> Babesia canis

<220>

<221> CDS

<222> (75) .. (500)

<400> 1

gaattcggca cgagccctgc tatactgtgc ttgcaacta actccatcgt aataatttaa 60

tataataata aagg atg gag tcg aca tca aca acg acc aac ttt gtt gcc 110

Met Glu Ser Thr Ser Thr Thr Thr Asn Phe Val Ala

1

5

10

gag aac cgt ccc acc ttt ggt gag acg ttt gat gtg atg agg gaa gct 158

Glu Asn Arg Pro Thr Phe Gly Glu Thr Phe Asp Val Met Arg Glu Ala

15

20

25

ttg ctt cgt gta aag tcc tct gaa cgc ttg gca atg ctc aga gcg ctt 206

Leu Leu Arg Val Lys Ser Ser Glu Arg Leu Ala Met Leu Arg Ala Leu

30

35

40

gca gga atg tgc ggt cac cgc gtc ctt cct ggc act ggt gct tct gcg 254

Ala Gly Met Cys Gly His Arg Val Leu Pro Gly Thr Gly Ala Ser Ala

45

50

55

60

ata gcg gca acg gta acc cca aag ggg gct tcg atg aag ctt aaa cca 302

gag atg aag cag tca aag aag tct cag aag ctc aag gag tct tca aag

Pro Arg Pro Gln Ser Thr Lys Ser Pro Glu Leu Arg Glu Leu Ser Arg

80

85

90

aag att cgc gaa atg aat aag act ata agt cag gaa tca gct cgg gta 398

Lys Ile Arg Glu Met Asn Lys Thr Ile Ser Gln Glu Ser Ala Arg Val

95

100

105

aac cac cgg ttg ccg gaa ggc cac cct ctc tta gag aag cgg gca gaa 446

Asn His Arg Leu Pro Glu Gly His Pro Leu Leu Glu Lys Arg Ala Glu

110

115

120

tat ttt cgt cac ctt aga tct ctt aag agc caa gga gtc aat aga ctc 494

Tyr Phe Arg His Leu Arg Ser Leu Lys Ser Gln Gly Val Asn Arg Leu

125

130

135

140

atc taa gaaggcacta cgtaggtagc gtgcctctat gaggaatacgc aaccgactag 550

Ile

tgcacaatag acgaccagtt ctaccaaagg tagagcctga ctctaatacta ccattcggcc 610

agcgacggag tcgcatgaca acgtggaatc ttagaccacg ccggacgggt tatccgtcaa 670

atggtacttt ggcagttacg gaactcctga tctcgattta tagatcaaac ttctacacct 730

tgaaggtggt cgaggaaggg agatgtacgt gctgcaacac ccataaggag caagctttgc 790

tactcctatc cggttacctc cagctatatc gtgcactgca ctgagttgga aggtctgtat 850

tcgtagaata ctgcaaaacc aggatatgcg tcgaggcacg cctcaccgga ctacgtccga 910

gggtgaccct aacgggctgc tgaactaggt tcagccagcg ctctcctgtga gtatgtcatt 970

ccgggtcctt cggggcccg gccagtttcg actggtgtag gtttgcccta ctagagtact 1030

tgcgacgccg aagcgctcc gttcaaaaga acgcaagc cctagcagag aaatgcgagg 1090

gcatgactct tcgagtcaaa aaaaaaaaaa aaaaaaaac tcgag 1135

<210> 2

<211> 141

<212> PRT

<213> Babesia canis

<400> 2

Met Glu Ser Thr Ser Thr Thr Thr Asn Phe Val Ala Glu Asn Arg Pro

1

5

10

15

Thr Phe Gly Glu Thr Phe Asp Val Met Arg Glu Ala Leu Leu Arg Val
 20 25 30

Lys Ser Ser Glu Arg Leu Ala Met Leu Arg Ala Leu Ala Gly Met Cys
 35 40 45

Gly His Arg Val Leu Pro Gly Thr Gly Ala Ser Ala Ile Ala Ala Thr
 50 55 60

Val Thr Pro Lys Gly Ala Ser Met Lys Leu Lys Pro Pro Arg Pro Gln
 65 70 75 80

Ser Thr Lys Ser Pro Glu Leu Arg Glu Leu Ser Arg Lys Ile Arg Glu
 85 90 95

Met Asn Lys Thr Ile Ser Gln Glu Ser Ala Arg Val Asn His Arg Leu
 100 105 110

Pro Glu Gly His Pro Leu Leu Glu Lys Arg Ala Glu Tyr Phe Arg His
 115 120 125

Leu Arg Ser Leu Lys Ser Gln Gly Val Asn Arg Leu Ile
 130 135 140

<210> 3

<211> 1134

<212> DNA

<213> Babesia canis

<220>

<221> CDS

<222> (75)..(929)

<400> 3

gaattcggca cgagccctgc tatactgtgc ttgcaacta actccatcgt aataatttaa 60

tataataata aagg atg gag tcg aca tca aca acg acc aac ttt gtt gcc 110
 Met Glu Ser Thr Ser Thr Thr Thr Asn Phe Val Ala
 1 5 10

gag aac cgt ccc acc ttt ggt gag acg ttt gat gtg atg agg gaa gct 158
 Glu Asn Arg Pro Thr Phe Gly Glu Thr Phe Asp Val Met Arg Glu Ala

Leu Leu Arg Val Lys Ser Ser Glu Arg Leu Ala Met Leu Arg Ala Leu

gca gga atg tgc ggt cac cgc gtc ctt cct ggc act ggt gct tct gcg 254
 Ala Gly Met Cys Gly His Arg Val Leu Pro Gly Thr Gly Ala Ser Ala
 45 50 55 60

ata gcg gca acg gta acc cca aag ggg gct tcg atg aag ctt aaa cca 302
 Ile Ala Ala Thr Val Thr Pro Lys Gly Ala Ser Met Lys Leu Lys Pro
 65 70 75

ccg cgt ccg cag tca acg aag tct ccg gag ctc agg gag ctg tca cgg 350
 Pro Arg Pro Gln Ser Thr Lys Ser Pro Glu Leu Arg Glu Leu Ser Arg
 80 85 90

aag att cgc gaa atg aat aag act ata agt cag gaa tca gct cgg gta 398
 Lys Ile Arg Glu Met Asn Lys Thr Ile Ser Gln Glu Ser Ala Arg Val
 95 100 105

aac cac cgg ttg ccg gaa ggc cac cct ctc tta gag aag cgg gca gaa 446
 Asn His Arg Leu Pro Glu Gly His Pro Leu Leu Glu Lys Arg Ala Glu
 110 115 120

tat ttc gtc acc tta gat ctc tta aga gcc aag gag tca ata gac tca 494
 Tyr Phe Val Thr Leu Asp Leu Leu Arg Ala Lys Glu Ser Ile Asp Ser
 125 130 135 140

tct aag aag gca cta cgt agg tac cgt gcc tct atg agg aat acg aac 542
 Ser Lys Lys Ala Leu Arg Arg Tyr Arg Ala Ser Met Arg Asn Thr Asn
 145 150 155

cga cta gtg cac aat aga cga cca gtt cta cca aag gta gag cct gac 590
 Arg Leu Val His Asn Arg Arg Pro Val Leu Pro Lys Val Glu Pro Asp
 160 165 170

tct aat cta cca ttc ggc cag cga cgg agt cgc atg aca acg tgg aat 638
 Ser Asn Leu Pro Phe Gly Gln Arg Arg Ser Arg Met Thr Thr Trp Asn
 175 180 185

ctt aga cca cgc cgg acg ggt tat ccg tca aat ggt act ttg gca gtt 686
 Leu Arg Pro Arg Arg Thr Gly Tyr Pro Ser Asn Gly Thr Leu Ala Val
 190 195 200

acg gaa ctc ctg atc tcg att tat aga tca aac ttc tac acc ttg aag 734
 Thr Glu Leu Leu Ile Ser Ile Tyr Arg Ser Asn Phe Tyr Thr Leu Lys
 205 210 215 220

gtg gtc gag gaa ggg aga tgt acg tgc tgc aac acc cat aag gag caa 782
 Val Val Glu Glu Gly Arg Cys Thr Cys Cys Asn Thr His Lys Glu Gln

225

230

235

gct ttg cta ctc cta tcc ggt tac ctc cag cta tat cgt gca ctg cac 830
 Ala Leu Leu Leu Leu Ser Gly Tyr Leu Gln Leu Tyr Arg Ala Leu His
 240 245 250

tca gtt gga agg tct gta ttc gta gaa tac tgc aaa acc agg ata tgc 878
 Ser Val Gly Arg Ser Val Phe Val Glu Tyr Cys Lys Thr Arg Ile Cys
 255 260 265

gtc gag gca cgc ctc acc gga cta cgt ccg agg gtg acc cta acg ggc 926
 Val Glu Ala Arg Leu Thr Gly Leu Arg Pro Arg Val Thr Leu Thr Gly
 270 275 280

tgc tgaactaggt tcagccagcg cttcctgtga gtatgtcatt ccgggtcctt 979
 Cys
 285

cggggcccgg gccagtttcg actggtgtag gtttgcccta ctagagtact tgcgacgccg 1039

aagcgccctcc gttcaaaaga acgcgcaagc cctagcagag aaatgcgagg gcatgactct 1099

tcgagtcaaa aaaaaaaaaa aaaaaaaaaac tcgag 1134

<210> 4

<211> 285

<212> PRT

<213> Babesia canis

<400> 4

Met Glu Ser Thr Ser Thr Thr Thr Asn Phe Val Ala Glu Asn Arg Pro
 1 5 10 15

Thr Phe Gly Glu Thr Phe Asp Val Met Arg Glu Ala Leu Leu Arg Val
 20 25 30

Lys Ser Ser Glu Arg Leu Ala Met Leu Arg Ala Leu Ala Gly Met Cys
 35 40 45

Gly His Arg Val Leu Pro Gly Thr Gly Ala Ser Ala Ile Ala Ala Thr
 50 55 60

Val Thr Pro Lys Gly Ala Ser Met Lys Leu Lys Pro Pro Arg Pro Gln

Met Asn Lys Thr Ile Ser Gln Glu Ser Ala Arg Val Asn His Arg Leu
 100 105 110
 Pro Glu Gly His Pro Leu Leu Glu Lys Arg Ala Glu Tyr Phe Val Thr
 115 120 125
 Leu Asp Leu Leu Arg Ala Lys Glu Ser Ile Asp Ser Ser Lys Lys Ala
 130 135 140
 Leu Arg Arg Tyr Arg Ala Ser Met Arg Asn Thr Asn Arg Leu Val His
 145 150 155 160
 Asn Arg Arg Pro Val Leu Pro Lys Val Glu Pro Asp Ser Asn Leu Pro
 165 170 175
 Phe Gly Gln Arg Arg Ser Arg Met Thr Thr Trp Asn Leu Arg Pro Arg
 180 185 190
 Arg Thr Gly Tyr Pro Ser Asn Gly Thr Leu Ala Val Thr Glu Leu Leu
 195 200 205
 Ile Ser Ile Tyr Arg Ser Asn Phe Tyr Thr Leu Lys Val Val Glu Glu
 210 215 220
 Gly Arg Cys Thr Cys Cys Asn Thr His Lys Glu Gln Ala Leu Leu Leu
 225 230 235 240
 Leu Ser Gly Tyr Leu Gln Leu Tyr Arg Ala Leu His Ser Val Gly Arg
 245 250 255
 Ser Val Phe Val Glu Tyr Cys Lys Thr Arg Ile Cys Val Glu Ala Arg
 260 265 270
 Leu Thr Gly Leu Arg Pro Arg Val Thr Leu Thr Gly Cys
 275 280 285

<210> 5
 <211> 90
 <212> DNA
 <213> Babesia canis

<400> 5
 ggatcctaatacgaactcactatagggagaccacatggagtcgacatcaa caacgaccaa 60
 ctttggtgccgagaaccgtccacctttgg 90

<210> 6	
<211> 24	
<212> DNA	
<213> Babesia canis	
<400> 6	
gacgtttgat gtgatgaggg aagc	24
<210> 7	
<211> 21	
<212> DNA	
<213> Babesia canis	
<400> 7	
aatgacatac tcacaggaag c	21
<210> 8	
<211> 20	
<212> DNA	
<213> Babesia canis	
<400> 8	
atgagtctat tgactccttg	20
<210> 9	
<211> 21	
<212> DNA	
<213> Babesia canis	
<400> 9	
agggagctgt cacggaagat t	21
<210> 10	
<211> 21	
<212> DNA	
<213> Babesia canis	
<400> 10	
atqagqaatt cgaaccgact a	21

